

Homework 8 — MATH 2510 Spring 2018

Consider the vocabulary $\mathcal{V} = \{\preceq\}$. Recall the following \mathcal{V} -sentences (called “axioms”) from Tuesday’s class:

1. **Axiom 1** (reflexive) $\forall x(x \preceq x)$
2. **Axiom 2** (anti-symmetric). $\forall x \forall y(x \preceq y \wedge y \preceq x \rightarrow x = y)$
3. **Axiom 3** (transitive). $\forall x \forall y \forall z(x \preceq y \wedge y \preceq z \rightarrow x \preceq z)$
4. **Axiom 4** (total). $\forall x \forall y(x \preceq y \vee y \preceq x)$

And recall the “well-ordering principle”: **every** nonempty subset of the universe contains a least element. (*note: writing this in logic symbols necessitates something like “ $\forall S \subseteq U$ ” which is not part of our language of first order logic — it is “second order logic”!*)

We say that a structure $M = (S | \preceq)$ is a preorder on S if it is reflexive and transitive. We say that a structure $M = (S | \preceq)$ is a partial order on S if it is an anti-symmetric preorder. We say that a structure $M = (S | \preceq)$ is a total order on S if it is a total partial order. We say that a structure $M = (S | \preceq)$ is a well-order on S if it is a total order that also obeys the well-ordering principle.

Problems

In every problem, determine which of the axioms that the structure models. If the structure is a preorder, partial order, total order, or well-order, then state so.

1. Consider the structure $M = (\{\emptyset, \{a\}, \{b\}, \{a, b\}\} | \preceq)$ where $x \preceq y$ is interpreted to mean $x \subseteq y$.
2. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be the function $f(x) = x^2$. Consider the structure $M = (\mathbb{R} | \preceq)$, where $x \preceq y$ is interpreted to mean $f(x) \leq f(y)$ (where “ \leq ” here is interpreted in the usual way).
3. Consider the xy -plane $\mathbb{R}^2 = \{(x, y) : x \in \mathbb{R}, y \in \mathbb{R}\}$. Consider the structure $M = (\mathbb{R}^2 | \preceq)$ where $(x_1, y_1) \preceq (x_2, y_2)$ is interpreted to mean that $x_1 \leq x_2$ and $y_1 \leq y_2$ (where “ \leq ” here is interpreted in the usual way).
(*note: this is called the product order of \leq*)
4. Consider the xy -plane \mathbb{R}^2 . Consider the structure $M = (\mathbb{R}^2 | \preceq)$ where $(x_1, y_1) \preceq (x_2, y_2)$ is interpreted to mean that $x_1 < x_2$ OR $x_1 = x_2$ and $y_1 \leq y_2$ (where “ $<$ ” and “ \leq ” are interpreted in the usual way). So for example, $(-2, 5) \preceq (5, 2000)$ because $-2 < 5$ and $(-2, 232) \preceq (-2, 333)$ because $-2 = -2$ and $232 \leq 333$.
(*note: this is called “lexicographic” or “dictionary” order*)

5. (Sharkovsky order) Consider the structure $M = (\mathbb{N} | \preceq)$, where \preceq is interpreted to be the so-called Sharkovskii order:

$$1 \prec 2 \prec 2^2 \prec 2^3 \prec \dots \prec 2^n \prec \dots \quad \dots \prec 7 \cdot 2^n \prec 5 \cdot 2^n \prec 3 \cdot 2^n \prec \dots \\ \dots \prec 7 \cdot 2 \prec 5 \cdot 2 \prec 3 \cdot 2 \prec \dots \prec 9 \prec 7 \prec 5 \prec 3.$$

You can understand this interpretation by thinking of it as saying

$$\text{descending powers of 2 (ascending)} \preceq \dots \preceq (\text{odd numbers}) \cdot 2^n \preceq (\text{odd numbers}) \cdot 2^{n-1} \preceq \dots \preceq (\text{odd numbers}) \cdot 2 \preceq \text{odd numbers (descending)}$$

(note: this order is fundamental to Sharkovskii's theorem from the 1960's)

Recall that we defined the notation $A \subseteq B$ to abbreviate the formula $z \in A \rightarrow z \in B$. We write $A \cap B$ as an abbreviation for $z \in A \wedge z \in B$.

6. Prove that that $X \cap Y \subseteq X$.