

International Workshop on Nonlinear Dynamical Systems and Functional Analysis

August 13-16, 2018

Universidade de Brasília

ICM 2018 Satellite Conference



Organizing Committee:

Carlos Alberto dos Santos (U. de Brasília/Brazil)
Elves Alves Barros e Silva (U. de Brasília/Brazil)
Everaldo de Mello Bonotto (U. de São Paulo/Brazil)
Giovany Figueiredo (U. de Brasília/Brazil)
Jaqueline Godoy Mesquita (U. de Brasília/Brazil)
Jiazheng Zhou (U. de Brasília/Brazil)
Juliana Pimentel (U. Federal do ABC/Brazil)
Liliane de Almeida Maia (U. de Brasília/Brazil)
Luís Henrique de Miranda (U. de Brasília/Brazil)
Manuela Rezende (U. de Brasília/Brazil)
Marcelo Furtado (U. de Brasília/Brazil)
Ricardo Parreira da Silva (U. de Brasília/Brazil)
Ricardo Ruviano (U. de Brasília/Brazil)

Scientific Committee:

Alexandre N. de Carvalho (U. de São Paulo/Brazil)
Carlos Lizama (U. de Santiago de Chile/Chile)
Hans-Otto Walther (U. of Giessen/Germany)
Jean Mawhim (U. Catholique de Louvain/Belgium)
João Marcos Bezerra do Ó (U. de Brasília/Brazil)
José Bonet (U. Politécnica de Valencia/Spain)
Pedro Miana (U. de Zaragoza/Spain)

Sponsors:



UnB



Fundação de Empreendedorismo Científico e Tecnológico



Fundação de Amparo à Pesquisa do Distrito Federal



Access homepage
of the event using
this QR code

INTERNATIONAL WORKSHOP ON NONLINEAR DYNAMICAL SYSTEMS AND FUNCTIONAL ANALYSIS

Book of Abstracts

Scientific Committee

Alexandre Nolasco de Carvalho (Universidade de São Paulo)
Carlos Lizama (Universidad de Santiago de Chile)
Hans-Otto Walther (University of Giessen)
Jean Mawhin (Université Catholique de Louvain)
João Marcos Bezerra do Ó (Universidade de Brasília)
José Bonet (Universidad Politécnica de Valencia)
Pedro Miana (Universidad de Zaragoza)

Organizing Committee

Carlos Alberto dos Santos (Universidade de Brasília)
Elves Alves Barros e Silva (Universidade de Brasília)
Everaldo de Mello Bonotto (Universidade de São Paulo)
Giovany Figueiredo (Universidade de Brasília)
Jaqueline Mesquita (Universidade de Brasília)
Jiazheng Zhou (Universidade de Brasília)
Juliana Pimentel (Universidade Federal do ABC)
Liliane de Almeida Maia (Universidade de Brasília)
Luís Henrique de Miranda (Universidade de Brasília)
Manuela Rezende (Universidade de Brasília)
Marcelo Furtado (Universidade de Brasília)
Ricardo Parreira (Universidade de Brasília)
Ricardo Ruviano (Universidade de Brasília)

Local Committee

Aldo Pereira Solis (Universidade de Brasília)
Anna Carolina Lafetá (Universidade de Brasília)
Gustavo Silvestre do Amaral Costa (Universidade de Brasília)
Karla Carolina Vicente de Sousa (Universidade de Brasília)
Laís Moreira dos Santos (Universidade de Brasília)
Mateus Malta Fleury (Universidade de Brasília)
Mayra Soares Costa Rodrigues (Universidade de Brasília)
Ricardo Lima Alves (Universidade de Brasília)

Session Organizers

Difference Equations/Dynamic equations on time scales

Organizer: Eduard Toon (Universidade Federal de Juiz de Fora)

Elliptic Partial Differential Equations

Organizer: Edcarlos Domingos da Silva (Universidade Federal de Goiás)

Functional Analysis

Organizer: Geraldo de Azevedo Botelho (Universidade Federal de Uberlândia)

Nonlinear Dynamical Systems

Organizer: Juliana Pimentel (Universidade Federal do ABC)

Ordinary/Functional Differential Equations

Organizer: Pierluigi Benevieri (Universidade de São Paulo)

Parabolic Differential Equations

Organizer: Anderson Luis Albuquerque de Araújo (Universidade Federal de Viçosa)


Poster

Organizer: Willian Cintra (Universidade de Brasília)

Adress

Universidade de Brasília
Instituto de Ciências Exatas
Departamento de Matemática
Campus Universitário Darcy Ribeiro
70910-900 Brasília - DF

Map of the University of Brasília



1- HUB (Hospital Universitário) - University hospital	16- MASC	31- Centro Comunitário Athos Bulcão - Community Center
2- Finatec	17- BSA	32- CIC
3- FUBRA	18- Instituto de Biologia - Biology Institute	33- UED
4- AUTOTRAC	19- PMU II	34- Almoarifado - Warehouse
5- FIO CRUZ	20- PMU II	35- PIC
6- CAEP	21- FE	36- PAT
7- CET	22- Casa do Professor	37- MASC
8- CRAD	23- SG 1 ao SG 12	38- FACE
9- CDT	24- Banco do Brasil - Bank	39- FA
10- CIME	25- Restaurante Universitário - University Restaurant	40- ASFUB
11- NIMT	26- ICC	41- Posto de Gasolina/Subway/Spoletto - Gas Station/Restaurant
12- CPD	27- Departamento de Matemática - Math department	42- PMDF - Police
13- CESPE	28- Teatro de Arena Honestino Guimarães - Arena Theater	43- Colina
14- Faculdade de Saúde - Health College	29- Reitoria - Rectory	44- FT
15- Instituto de Química - Chemistry Institute	30- Biblioteca - Library	45- Auditório da FT

Contents

General Information	11
Location	12
Important Numbers	12
Meals and refreshments	12
Schedule	13
Plenary Talks	16
Alexandre Nolasco	
Non-Autonomous Morse-Smale Dynamical Systems: Structural Stability under Non-Autonomous Perturbations	17
Carlos Rocha	
Design of dynamic complexes for Sturm global attractors	18
Gabriela Planas	
On the non-isothermal Navier-Stokes-Allen-Cahn equations for two-phase fluids	19
Gennaro Infante	
Positive solutions of nonlocal elliptic systems with functional BCs	20
Hans-Otto Walther	
Differential Equations with State-Dependent Delay, Fréchet and other Derivatives, and Complicated Motion with functional BCs	21
José Galé	
On reproducing kernels and fractional spaces.	22
Mahamadi Warma	
Optimal control of fractional semilinear PDEs	23
Márcia Federson	
On the Schrödinger equation in the quantum case	24
Martin Bohner	
Massera's Theorem in Quantum Calculus	25
Olimpio H. Miyagaki	
Hénon type problems: nonradial solutions, asymptotic behavior, compactness, etc.	26
Pedro J. Miana	
Evolution C_0 -semigroups and generalized Cesáro operators	27
Tibor Krisztin	
A differential equation with a state-dependent queueing delay	28
Special Sessions	29
Difference Equations/Dynamic Equations on time scales	30
Aldo Pereira	
Cosine and sine functions on time scales	31
Eduard Toon	
Boundedness of solutions of dynamic equations on time scales	32
Fernanda Andrade da Silva	
Controllability for dynamic systems on time scales	33
Geraldo Silva	
Optimal Control on Time Scales: Necessary Optimality Conditions	34

Sabrina Streipert	
Periodic functions on isolated Time Scales	35
Tom Cuchta	
Lambert W for time scales	36
Ordinary/Functional Differential Equations	37
Agnieszka Malinowska	
On systems of fractional differential equations with a general derivative	38
Carlos Tomei	
Beyond integrability: real life eigenvalue algorithms	39
João Marcos do Ó	
Elliptic equations and systems with critical Trudinger-Moser nonlinearities	40
Maria Carolina Mesquita	
Bifurcation theory for generalized ODE's	41
Marielle Silva	
On oscillation of Functional Differential Equations with delay with values in \mathbb{R}^n	42
Pablo Amster	
Uniform persistence and periodic solutions for some systems of delay differential equations	43
Pierluigi Benevieri	
Global persistence of nonlinear operators in Hilbert spaces	44
Valeriano Antunes	
Optimality Conditions for Optimal Control Problems with Mixed Constraints	45
Nonlinear Dynamical Systems	47
André Caldas	
The Variational Principle for Locally Compact Spaces	48
Everaldo de Mello Bonotto	
Impulsive non-autonomous dynamical systems	49
Jacson Simsen	
Construction of a global solution for a process	50
Juliana Berbert	
Combining random searches, individual's memory, and population dynamics: analysis of population redistribution patterns	51
Lucas Seco	
The minimal Morse components of translations in flag manifolds are normally hyperbolic	52
Mauro Patrão	
The Topological Entropy of Endomorphisms of Lie Groups	53
Marcos C. Pereira	
A nonlocal approach to spatial spread in thin structures	54
Phillipo Lappicy	
Unbounded Sturm attractors for quasilinearparabolic equations	55
Piotr Kalita	
Rayleigh–Bénard problem for micropolar fluids	56
Rodrigo A. Samprognna	
Robustness with respect to exponents for non-autonomous reaction diffusion equations	57
Sabrina Camargo	
Intermingled basins in coupled Lorenz systems	59

Functional Analysis	60
Alex F. Pereira	
Duality in Spaces of Lorch Analytic Mappings	61
Claudia Correa	
Twisted sums of c_0 and $C(K)$	62
Daniela M. Vieira	
On the spectrum of the algebra of bounded type holomorphic functions on Banach spaces	63
Elisa R. Santos	
Some stability properties of the polynomial daugavetian index	64
Gustavo Araújo	
Summability of multilinear operators: a unified theory and consequences	65
Geraldo Botelho	
Jorge Mujica's last theorem and applications	66
Jamilson R. Campos	
mid (q, p) -summable sequences	67
Khazhak Navoyan	
Complementation in Fremlin vector lattice symmetric tensor products	68
M. Lilian Lourenço	
Biholomorphic Mappings on Banach Spaces	69
Nilson C. Bernardes Jr.	
Mean Li-Yorke Chaotic Operators on Banach Spaces	70
Santiago Muro	
Dynamics of homogeneous polynomials on Banach spaces	71
Thiago R. Alves	
Holomorphic functions with distinguished properties on infinite dimensional spaces . .	72
Victor Simões Barbosa	
Positive definite kernels on two-point homogeneous spaces	73
Elliptic Partial Differential Equations	74
Damião J. Araújo	
Sharp Regularity for the Inhomogenous Porous Medium Equation	75
Diego Moreira	
Up to the boundary gradient estimates in free boundary problems	76
Eduardo Domingos	
On nonquadratic fractional coupled elliptic systems in unbounded domains	77
Eudes Mendes Barboza	
Hénon Elliptic Equations in \mathbb{R}^2 with Subcritical and Critical Exponential Growth: Mountain–Pass case	78
Gustavo Madeira	
Infinitely many solutions for a class of degenerated elliptic p-Kirchhoff equations . . .	81
Gaetano Siciliano	
Least energy nodal solution for the Schrödinger-Poisson system under an asymptotically cubic nonlinearity	82
José Francisco de Oliveira	
Extremal for a k -Hessian inequality of Trudinger-Moser type	83
Jorge Marques	
On the well-posedness of Goursat problems in Gevrey classes	84

Kaye Oliveira Silva	
The fibering method approach for a non-linear Schrödinger equation coupled with the electromagnetic field	85
Maxwell Lizete	
On the well-posedness of Goursat problems in Gevrey classes	86
Minbo Yang	
On the Choquard equation with Hardy-Littlewood-Sobolev upper critical exponent . .	87
Pawan Mishra	
Super critical problems with concave and convex nonlinearities in \mathbb{R}^N	88
Parabolic Differential Equations	89
Adilson Presoto	
The number π and the nonexistence of solution for Chern-Simons equation	90
Anderson L.A. de Araujo	
A class of parabolic equations driven by the mean curvature flow	91
Arlúcio Viana	
Global solutions for a fractional reaction–diffusion equation	92
Jamil Abreu	
On the Dirichlet-to-Neumann Semigroup: Recent Developments and Results	93
Luis Henrique de Miranda	
Fractional Regularity for Degenerate Equations	94
Rodrigo Nunes Monteiro	
Attractors for a vectorial von Karman system	95
Poster Session	96
Alfredo Sotelo Pejerrey	
Elliptic equations and systems with critical Trudinger-Moser nonlinearities	97
Angelina Carrijo	
Relation between approximation tools in homogeneous compact spaces of rank 1 . . .	98
Cláudia Santana	
Standing waves for a system of quasilinear schrödinger equations in \mathbb{R}^N	99
Claudiney Goulart	
Standing waves for weakly coupled nonlinear Schrodinger systems with critical growth in \mathbb{R}^N	101
Daniel Raom Santiago	
Mountain pass algorithm via Pohozaev manifold	102
Felipe B. Federson	
On retarded potentials and applications	103
Fernando S. Carvalho	
Topological optimization in the kirchhoff plate bending model using topological derivative	104
Ian Rodrigues Massa	
Solving D’Alambert problem using Perplex Numbers	105
José Carlos de Albuquerque	
Positive ground states for a subcritical and critical coupled system involving Kirchhoff-Schrödinger equations	106
Lais Santos	
<i>Continuums</i> of positive solutions for classes of non-autonomous and non-local problems with strong singular term	107

Leonardo Epiphonio Galvão	
Singularity formation in toy models for incompressible flow	108
Marcos L. M. Carvalho	
A Brézis-Oswald problem to Φ -Laplacian operator with a gradient term	109
Mateus Malta Fleury	
Existence and uniqueness of solutions for RFDEs	110
Mayra S. C. Rodrigues	
Asymptotically Linear Indefinite Problems in \mathbb{R}^N via an Abstract Linking Theorem	111
Monisse Postigo Alves	
Existence of solutions in Besov-Morrey spaces for chemotaxis Navier-Stokes fluid with chemical attractant	112
Paulo N. Seminario Huertas	
Global well-posedness of weak solutions for the Vlasov-Fokker-Planck system	113
Pedro Gabriel Ferreira Colucci	
On the Quantum Harmonic Oscillator Model applied to Spectroscopy	114
Ricardo Lima Alves	
Existence of multiple positive solutions for Schrödinger equation with singular nonlinearities sign-changing	115
Rodrigo Clemente	
On Lane-Emden systems with singular nonlinearities	116
Sanjeev Kumar	
A model for determining the concentration of urea in artificial kidney	117
Steffânio Moreno de Sousa	
Sistemas com Termo Cncavo-Convexo em Domnio No Limitado	118
Thiago Rodrigues Cavalcante	
Fourth order elliptic problems with combined nonlinearities	119

Sponsors **120**

General Information

Location

The congress will take place in FINATEC at the University of Brasília. Also, the participant may want to know the department of mathematics of the University of Brasília. Both places are indicated in blue in the map on page 5.

Useful Phone Numbers

In case of any health emergencies call 192 (SAMU).

Police number: 190.

Math department of the University of Brasília: (61) 3107 7236

Security of the University of Brasília: (61) 3107 6222

Meals and refreshments

There is a university restaurant indicated in the map of page 4 that serves breakfast (7:00hs - 9:00hs), lunch (11:00hs - 14:30hs) and dinner (17:00hs - 19:30hs). There are several restaurants nearby the in Asa Norte. We will present you a few options:

1. Feitiço Mineiro located at 306 Norte, bloco B - lojas 45/51, Asa Norte (lunch and dinner).
2. Restaurante e Bar Xique Xique located at 708 Norte Bloco E Loja 45, Asa Norte (lunch and dinner).
3. Subway/Spoletto at the University of Brasília, indicated in the map of page 4, number 41 (lunch and dinner).
4. Domino's Pizza located at 109 Norte Bloco B, Loja 1, Asa Norte (Dinner).
5. Crepe au Chocolat located at 109 Norte, Bloco C, Loja 5, Asa Norte (Launch and Dinner).
6. Restaurante El Negro located at 413 Norte Bloco C, Loja 21, Asa Norte.

There is also a supermarket (Pão de Açúcar) at 404/405 Norte, Bloco A, Asa Norte opened from 7:00hs to 22:00hs.

Schedule

International Workshop on Nonlinear Dynamical Systems and Functional Analysis

Auditorium	Plenary Talks			
	Monday 13	Tuesday 14	Wednesday 15	Thursday 16
Chairman	<i>Jaqueline Mesquita</i>	<i>Juliana Pimentel</i>	<i>Giovany Figueiredo</i>	<i>Luis Miranda</i>
09:00 - 09:40		Bernold Fiedler	Tibor Krisztin	Gennaro Infante
09:40 - 10:20	Hans-Otto Walther	Carlos Rocha	Pedro Miana	Márcia Federson
10:20 - 11:00	Coffee Break/Posters	Coffee Break/Posters	Coffee Break/Posters	Coffee Break
11:00 - 11:40	Martin Bohner	Alexandre Nolasco	Gabriela Planas	Minbo Yang
11:40 - 12:20	Carlos Tomei	Photo	Oimpio Miyagaki	
12:20 - 14:30	Lunch	Barbecue (13:00-18:00)	Lunch	Lunch

T.B.A.	Special Session on Parabolic Differential Equations			
	Monday 13	Tuesday 14	Wednesday 15	Thursday 16
Chairman	<i>Luis Miranda</i>	Barbecue	Coffee Break	Coffee Break
14:30 - 15:00	Adilson Presoto			
15:00 - 15:30	Anderson Araújo			
15:30 - 16:00	Jamil Abreu			
16:00 - 16:30	Coffee Break			
Chairman	<i>Anderson Araújo</i>			
16:30 - 17:00	Arlúcio Viana			
17:00 - 17:30	Luis Miranda			
17:30 - 18:00	Rodrigo Monteiro			Closing

T.B.A.	Special Session on Functional Analysis			
	Monday 13	Tuesday 14	Wednesday 15	Thursday 16
Chairman	<i>Geraldo Botelho</i>	Barbecue	<i>Nilson Bernardes</i>	Coffee Break
14:30 - 15:00	Mary Lourenço		Khazhak Navoyan	
15:00 - 15:30	Nilson Bernardes		Santiago Muro	
15:30 - 16:00	Elisa Santos		Daniele Vieira	
16:00 - 16:30	Coffee Break		Coffee Break	
Chairman	<i>Pedro Miana</i>		<i>Mary Lourenço</i>	
16:30 - 17:00	Gustavo Araújo		Alex Pereira	
17:00 - 17:30	Cláudia Correa		Thiago Alves	
17:30 - 18:00	Geraldo Botelho	Jamilson Campos	Closing	
18:00 - 18:30		Victor Barbosa		

T.B.A.	Special Session on Nonlinear Dynamical Systems				
	Monday 13	Tuesday 14	Wednesday 15	Thursday 16	
Chairman		Barbecue	<i>Sergio Oliva</i>	<i>Everaldo Bonotto</i>	
14:30 - 15:00			Everaldo Bonotto	Mauro Patrão	
15:00 - 15:30			Juliana Pimentel	Philipo Lappicy	
15:30 - 16:00			Jacson Simsen	Lucas Seco	
16:00 - 16:30			Coffee Break	Coffee Break	Coffee Break
Chairman			<i>Mauro Patrão</i>		
16:30 - 17:00			Sergio Oliva		
17:00 - 17:30			Sabrina Carmargo		
17:30 - 18:00		Rodrigo Samprogna			

International Workshop on Nonlinear Dynamical Systems and Functional Analysis

Special Session on Elliptic Partial Differential Equations					
T.B.A.	Monday 13	Tuesday 14	Wednesday 15	Thursday 16	
Chairman	Maxwell Silva	Barbecue	Edcarlos Domingos		
14:30 - 15:00	Gaetano Siciliano		Damião Araújo		
15:00 - 15:30	Gustavo Madeira		Pawan Mishra		
15:30 - 16:00	Kaye Silva		José Francisco Alves		
16:00 - 16:30	Coffee Break		Coffee Break		Coffee Break
Chairman	Kaye Silva		Damião Araújo		Closing
16:30 - 17:00	Eudes Barboza		Edcarlos Domingos		
17:00 - 17:30	Diego Moreira		Maxwell Silva		
17:30 - 18:00			Jorge Marques		

Special Session on Ordinary/Functional Differential Equations					
T.B.A.	Monday 13	Tuesday 14	Wednesday 15	Thursday 16	
Chairman	Pablo Amster	Barbecue	Pierluigi Benevieri		
14:30 - 15:00	Marielle Silva		João Marcos do Ó		
15:00 - 15:30	Pierluigi Benevieri		Agnieszka Malinowska		
15:30 - 16:00	Maria Carolina Mesquita		Pablo Amster		
16:00 - 16:30	Coffee Break		Coffee Break		Coffee Break
Chairman	Pablo Amster		Closing		
16:30 - 17:00	Valeriano Antunes				
17:00 - 17:30	Joelma Azevedo				
17:30 - 18:00					

Special Session on Difference Equations/Dynamic Equations on Time Scales						
T.B.A.	Monday 13	Tuesday 14	Wednesday 15	Thursday 16		
Chairman	Eduard Toon	Barbecue				
14:30 - 15:00	Geraldo Silva					
15:00 - 15:30	Fernanda Silva					
15:30 - 16:00	Tom Cuchta					
16:00 - 16:30	Coffee Break				Coffee Break	Coffee Break
Chairman	Geraldo Silva				Closing	
16:30 - 17:00	Eduard Toon					
17:00 - 17:30	Aldo Solis					
17:30 - 18:00	Sabrina Streipert					

Poster Session				
T.B.A.	Monday 13	Tuesday 14	Wednesday 15	Thursday 16
10:20 - 11:00	Lais Santos	Cláudia Santana	Daniel Santiago	
	Angelina Faria	Claudiney Goulart	Fernando Carvalho	
	Felipe Federson	José Carlos Albuquerque	Ian Massa	
	Monisse Alves	Alfredo Pejerrey	Leonardo Galvão	
	Pedro Gabriel Colucci	Marcos Leandro Carvalho	Mateus Fleury	
	Sanjeev Kumar	Mayra Rodrigues	Paulo Huertas	
	Thiago Cavalcante	Ricardo Alves	Steffânio Sousa	
		Rodrigo Clemente		

Auditorium	Monday 13	Tuesday 14	Wednesday 15	Thursday 16
08:20 - 09:00	Registration	Photo (11h40)		
09:00 - 09:40	Opening			
11:40				
13:00 - 18:00		Barbecue		Closing (17h30)

Plenary Talks

Non-Autonomous Morse-Smale Dynamical Systems: Structural Stability under Non-Autonomous Perturbations

Alexandre Nolasco
Instituto de Ciências Matemáticas e de Computação
Universidade de São Paulo, São Carlos, Brazil

Abstract

In this lecture we present our recent results on structural stability of gradient Morse-Smale Dynamical Systems under non-autonomous perturbations. To that end we introduce the notion of lifted invariant sets and give a characterization of the uniform attractor in terms of dynamical structures of a family of pullback attractors. This is a joint work with G. Raugel (Paris XI), J. Langa (U. Sevilla) and M. Bortolan (UFSC-Brazil).

Design of dynamic complexes for Sturm global attractors

Carlos Rocha

Partially supported by FCT/Portugal through project UID/MAT/04459/2013

e-mail: crocha@math.ist.utl.pt

Center for Mathematical Analysis, Geometry and Dynamical Systems

Department of Mathematics

University of Lisbon

Lisbon, Portugal

Abstract

We use the geometric and combinatorial characterization of global Sturm attractors to discuss the Thom-Smale dynamic complexes consisting of a single 3D ball resulting from scalar semilinear parabolic differential equations with Neumann boundary conditions. This is a joint work with B. Fiedler.

References

- [1] B. Fiedler and C. Rocha, Sturm 3-balls and global attractors 1: Thom-Smale complexes and meanders, to appear in *São Paulo J. Math. Sc.*, arXiv:1611.02003, 2016.
- [2] B. Fiedler and C. Rocha, Sturm 3-balls and global attractors 2: Design of Thom-Smale complexes, to appear in *J. Dyn. Diff. Eqns.*, arXiv:1704.00344, 2017.
- [3] B. Fiedler and C. Rocha, Sturm 3-balls and global attractors 3: Examples of Thom-Smale complexes, to appear in *Discr. Cont. Dyn. Sys.*, arXiv:1708.00690, 2018.

On the non-isothermal Navier-Stokes-Allen-Cahn equations for two-phase fluids

Gabriela Planas
Departamento de Matemática
IMECC - Unicamp
Campinas, Brazil

Juliana Honda Lopes
Departamento de Matemática
IMECC - Unicamp
Campinas, Brazil

Abstract

This talk is concerned with some non-isothermal diffuse-interface models which describe the motion of a mixture of two viscous incompressible fluids. These kind of models consist of modified Navier-Stokes equations coupled with a phase-field equation given by a convective Allen-Cahn equation, and energy transport equation for the temperature. It is discussed questions related to the well-posedness of the problem in the two and three dimensional cases. Moreover, regular and singular potentials for the phase-field equation are considered.

Positive solutions of nonlocal elliptic systems with functional BCs

Gennaro Infante

Partially supported by G.N.A.M.P.A. - INdAM (Italy)

e-mail: gennaro.infante@unical.it

Dipartimento di Matematica e Informatica

Università della Calabria

Cosenza, Italy

Abstract

We discuss the existence of non-negative weak solutions for second order nonlocal elliptic systems subject to functional boundary conditions and depending on some parameters. This setting for the boundary conditions is fairly general and covers the case of multi-point, integral and nonlinear boundary conditions. Our approach is based on classical fixed point index theory and we make use of recent results contained in [1].

References

- [1] G. Infante, Nonzero positive solutions of a multi-parameter elliptic system with functional BCs, *Topol. Methods Nonlinear Anal.*, to appear.

Differential Equations with State-Dependent Delay, Fréchet and other Derivatives, and Complicated Motion

Hans-Otto Walther
E-mail: Hans-Otto.Walther@math.uni-giessen.de
Mathematisches Institut
Universität Gießen
Gießen, Germany

Abstract

The lecture presents (1) recent results about delay differential equations in the general form

$$x'(t) = f(x_t)$$

which are designed for the application to equations with state-dependent delay, which may be unbounded. The state space which carries a nice semiflow is a Fréchet manifold, and the (weak) assumptions on f which are required for the theory (well-posedness of initial value problem, linearization, local invariant manifolds) have implications on the nature of differentiability of f as well as on the type of delay (which can be formulated as a property of the functional f).

We also (2) report about work on complicated solution behaviour which seems to be specific for delay differential equations and which is less local than chaotic motion in a thin "Cantor-dust".

References

- [1] H. O. Walther. *Semiflows for differential equations with locally bounded delay on solution manifolds in the space $C^1((-\infty, 0], \mathbb{R}^n)$* . DOI 10.12775/TMNA.2016.056, Topological Methods in Nonlinear Analysis, **48** (2016), 507-537.
- [2] H. O. Walther. *Local invariant manifolds for delay differential equations with state space in $C^1((-\infty, 0], \mathbb{R}^n)$* . DOI 10.14232/ejqtde.2016.1.85, Electronic Journal of the Qualitative Theory of Differential Equations, **No. 85** (2016), 1-29.
- [3] H. O. Walther. *Fréchet differentiability in Fréchet spaces, and differential equations with unbounded variable delay*. arXiv: 1801.09213, preprint, 2018, 45 pp.
- [4] H. O. Walther. *A delay differential equation with a solution whose shortened segments are dense*. Preprint, 2017, submitted, 30 pp.
- [5] H. O. Walther. *Maps which are continuously differentiable in the sense of Michal and Bastiani but not of Fréchet*. Preprint, 2017, submitted, 13 pp.

On reproducing kernels and fractional spaces.

Jos Gal
Universidad de Zaragoza
Zaragoza, Spain

Abstract

Certain spaces of absolutely continuous functions of fractional higher degree are considered. Such spaces admit Brownian versions as well as models formed by holomorphic functions on the half plane, with reproducing kernels accordingly. The definition involves Cesro operators. This talk is part of joint work with P. J. Miana and L. Snchez-Lajusticia.

Optimal control of fractional semilinear PDEs

Mahamadi Warma

Partially supported by the US Air Force Office of Scientific Research under the Award No:
FA9550-18-1-0242

e-mail: mahamadi.warma1@upr.edu, mjwarma@gmail.com

Department of Mathematics

University of Puerto Rico, Rio Piedras Campus
San Juan, USA

Abstract

In this talk we consider the optimal control of semilinear fractional PDEs with both spectral and integral fractional diffusion operators of order $2s$ with $s \in (0, 1)$. We first prove the boundedness of solutions to both semilinear fractional PDEs under minimal regularity assumptions on domain and data. We next introduce an optimal growth condition on the nonlinearity to show the Lipschitz continuity of the solution map for the semilinear elliptic equations with respect to the data. We further apply our ideas to show existence of solutions to optimal control problems with semilinear fractional equations as constraints. Under the standard assumptions on the nonlinearity (twice continuously differentiable) we derive the first and second order optimality conditions.

References

- [1] H. Antil, J. Pfefferer and M. Warma. *A note on semilinear fractional elliptic equation: analysis and discretization*. ESAIM Math. Model. Numer. Anal. **51** (2017), 2049–2067.
- [2] W. Arendt, A.F.M. ter Elst and M. Warma. *Fractional powers of sectorial operators via the Dirichlet-to-Neumann operator*. Comm. Partial Differential Equations **43** (2018), 1–24.
- [3] M. Warma, Mahamadi. *The fractional relative capacity and the fractional Laplacian with Neumann and Robin boundary conditions on open sets*. Potential Anal. **42** (2015), 499–547.

On the Schrödinger equation in the quantum case

Márcia Federson

Supported by FAPESP and CNPq.

Email: federson@icmc.usp.br

Instituto de Ciências Matemáticas e de Computação

Universidade de São Paulo,

São Carlos, Brazil

Abstract

We address some issues within the Schrödinger equation in the quantum case and the Feynman path integral and we give a quick overhaul of the main contributions of the theory of non-absolute integration in the sense of Ralph Henstock.

References

- [1] E.M. Bonotto; M. Federson; P. Muldowney, A Feynman-Kač solution to a random impulsive equation of Schrödinger type, *Real Anal. Exchange* 36(1), (2010/2011), 107-148.
- [2] R.P. Feynman, Space-time approach to non-relativistic quantum mechanics, *Rev. Modern Phys.* 20(2), (1948), 367-387.
- [3] R.P. Feynman; R.B. Leighton; M. Sands, *The Feynman Lectures on Physics*, Addison-Wesley, 1964.
- [4] P. Muldowney, Feynman's path integrals and Henstock's non-absolute integration. *J. Appl. Anal.* 6(1), (2000), 1-24.
- [5] P. Muldowney, *A Modern Theory of Random Variation - The Henstock Integral in Probability and Stochastic Calculus, with application to Option Pricing and Feynman Path Integrals*, John Wiley & Sons, Inc. 2012.
- [6] E.S. Nathanson, *Path integration with non-positive distributions and applications to the Schrödinger equation*. PhD Thesis, The University of Iowa. 2014. 113 pp.
- [7] E.S. Nathanson,; P.E.T. Jorgensen, A global solution to the Schrödinger equation: from Henstock to Feynman. *J. Math. Phys.* 56(9), (2015), 092102, 15 pp.
- [8] E.S. Nathanson,; P.E.T. Jorgensen, Trotter's limit formula for the Schrödinger equation with singular potential. *J. Math. Phys.* 58(12) (2017), 122101, 13 pp.

Massera's Theorem in Quantum Calculus

Martin Bohner

E-mail: bohner@mst.edu

Departement of Mathematics and Statistics

Missouri S&T

Rolla, MO 65409-0020, USA

Abstract

In this talk, we present versions of Massera's theorem for linear and nonlinear q -difference equations and give some examples to illustrate our results. This is joint work with Jaqueline Mesquita (University of Brasilia, Brazil) and will appear in the *Proceedings of the American Mathematical Society*.

References

- [1] Martin Bohner and Rotchana Chieochan. Floquet theory for q -difference equations. *Sarajevo J. Math.*, 8(21)(2):1–12, 2012.
- [2] Martin Bohner and Jaqueline Mesquita. Periodic averaging principle in quantum calculus. *J. Math. Anal. Appl.*, 435(2):1146–1159, 2016.
- [3] Martin Bohner and Jaqueline Mesquita. Massera's theorem in quantum calculus. *Proc. Amer. Math. Soc.*, 2018. To appear.
- [4] Martin Bohner and Sabrina Streipert. The second Cushing–Henson conjecture for the Beverton–Holt q -difference equation. *Opuscula Math.*, 37(6):795–819, 2017.

Hénon type problems: nonradial solutions, asymptotic behavior, compactness, etc.

O. H. Miyagaki

Partially supported by CNPQ and FAPEMIG

e-mail: ohmiyagaki@gmail.com

Department of Mathematics

Federal University of Juiz de Fora

Juiz de Fora, Brazil

Abstract

In this talk a Hénon type problem in a unit ball is discussed. By using minimization arguments on Nehari manifold is established that symmetry breaking appear when the problem is asymptotic to a some limit problem. Asymptotic behavior of the ground state solution is discussed, also, some compactness results are established.

The talk is related to the papers [1, 2, 3, 4, 5, 6].

References

- [1] D. G. Costa, O. H. Miyagaki, M. Squassina, J. Yang, *Asymptotics of ground states for Hénon type systems with fractional diffusion*, “Contributions to Nonlinear Elliptic Equations and Systems, Progress in Nonlinear Differential Equations and Their Applications 86, A Tribute to Djairo Guedes de Figueiredo on the Occasion of his 80th Birthday”(2015), 133-161.
- [2] R. B. Assuno, O. H. Miyagaki, G. A. Pereira, B. M. Rodrigues, *On a class of nonhomogeneous equations of Hénon-type: Symmetry breaking and non radial solutions*. Nonlinear Anal., **165** (2017), 102–120.
- [3] P.C. Carrião, D.G. de Figueiredo, O.H. Miyagaki, *Quasilinear elliptic equations of the Hénon-Type: Existence of non- radial solutions*, Comm.Contemporary Math. **5**(2009), 783–798.
- [4] D. G. de Figueiredo, E. Moreira dos Santos, O. H. Miyagaki, *Critical Hyperbolas and multiple symmetric solutions to some strongly coupled elliptic systems*, Advanced Nonlinear Studies, **13**(2013), 359-371.
- [5] D. G. de Figueiredo, E. Moreira dos Santos, O. H. Miyagaki, *Sobolev spaces of symmetric functions and applications*, Journal of Functional Analysis, **261**(2011), 3735-3770.
- [6] D. G. Costa, D. G. de Figueiredo, E. Moreira dos Santos, O. H. Miyagaki, *in preparation*

Evolution C_0 -semigroups and generalized Cesàro operators

Pedro J. Miana

Partially supported by Project MTM2016-77710-P, DGI-FEDER, of the MCYTTS and Project E-64,

D.G. Aragón, Universidad de Zaragoza, Spain

e-mail: pjmiana@unizar.es

Departamento de Matemáticas

Universidad de Zaragoza

Zaragoza, Spain

Abstract

In this talk, we present a complete spectral research of generalized Cesàro operators on Sobolev-Lebesgue sequence spaces. The main idea is to subordinate such operators to suitable C_0 -semigroups on these sequence spaces. We introduce that family of sequence spaces using the fractional finite differences and we prove some structural properties similar to classical Lebesgue sequence spaces. In order to show the main results about fractional finite differences, we state equalities involving sums of quotients of Euler's Gamma functions. Finally, we display some graphical representations of the spectra of generalized Cesàro operators. Main results of this talk are included in a joint paper with L. Abadias ([1]).

References

- [1] L. Abadias and P.J. Miana. *Generalized Cesàro operators, fractional finite differences and Gamma functions*. J. Funct. Anal., **274**, (2018), 1424–1465.

A differential equation with a state-dependent queueing delay

Tibor Krisztin

Partially supported by the Hungarian Scientific Research Fund (OTKA)

e-mail: krisztin@math.u-szeged.hu

Bolyai Institute

University of Szeged

Szeged, Hungary

Abstract

We consider a differential equation with a state-dependent delay motivated by a queueing process. The time delay is determined by an algebraic equation involving the length of the queue. For the length of the queue a discontinuous differential equation holds. We formulate an appropriate framework to study the problem, and show that the solutions define a Lipschitz continuous semiflow. Then, in the developed framework, we show that the system has a slowly oscillating periodic solution.

This is a joint work with István Balázs (Szeged, Hungary).

Special Sessions

Difference Equations/Dynamic Equations on time scales

Organizer: Eduard Toon (UFJF)

Cosine and sine functions on time scales

Aldo Pereira

e-mail: pereirasolis@gmail.com

Departament of Mathematics

University of Brasília

Brasília, Brazil

Abstract

Abstract cosine and sine functions defined on a Banach space are useful tools in the study of wide classes of abstract evolution equations. In this work, we introduce a definition of cosine and sine functions on time scales, which unify the continuous, discrete and cases which are between these ones. Our definition includes several types of time scales. For instance, real numbers set, integers numbers set, quantum scales, among others. We study the relationship between the cosine function on time scales and its infinitesimal generator, proving several properties concerning it. Also, we study the sine functions on time scales, presenting their main properties. Finally, we apply our theory to study the homogeneous and inhomogeneous abstract Cauchy problem on time scales in Banach spaces. Joint work with Jaqueline G. Mesquita (U. de Brasília) and Rodrigo Ponce (U. de Talca, Chile).

References

- [1] M. Bohner, A. Peterson: *Dynamic Equations on Time Scales: An Introduction with Applications*, Birkhäuser, Boston, 2001.
- [2] M. Bohner, A. Peterson: *Advances in Dynamic Equations on Time Scales*, Birkhäuser, Boston, 2003.
- [3] H.R. Henríquez, C. Lizama, J.G. Mesquita: *Semigroups on time scales and applications to abstract Cauchy problems*, submitted.

Boundedness of solutions of dynamic equations on time scales

Eduard Toon

e-mail: `eduard.toon@ufjf.edu.br`

Departament of Mathematics
Universidade Federal de Juiz de Fora
Juiz de Fora, Brazil.

Abstract

This is a joint work with Márcia Federson, Rogélio Grau and Jaqueline G. Mesquita. We prove results concerning boundedness of solutions for dynamic equations on time scales. In order to prove our main results, we prove results about boundedness of solutions for generalized ordinary differential equations and boundedness of solutions for measure differential equations and we use the fact that dynamic equations on time scales represent a particular case of measure differential equations.

References

- [1] M. Bohner and A. Peterson, *Dynamic Equations on Time Scales: An Introduction with Applications*, Birkhäuser, Boston, 2001.
- [2] M. Federson, R. Grau, J. G. Mesquita and E. Toon, Boundedness of solutions of measure differential equations and dynamic equations on time scales *Journal of Differential Equations* 263 (1), 26-56, 2017
- [3] Š. Schwabik, *Generalized Ordinary Differential Equations*, World Scientific, Series in Real Anal., vol. 5, 1992.
- [4] A. Slavík, Dynamic equations on time scales and generalized ordinary differential equations, *Journal of Mathematical Analysis and Applications* 385, 534-550, 2012.

Controllability for dynamic systems on time scales

Fernanda Andrade da Silva
 Partially supported by CAPES
 e-mail: ffeandrade@usp.br
 Marcia Federson; (e-mail: federson@icmc.usp.br)
 Eduard Toon (e-mail: edutoon@gmail.com)
 Departament of Mathematics
 University of São Paulo
 São Carlos, Brazil.

Abstract

In the literature, we found results on controllability for the following dynamic equation on times scales

$$x^\Delta(t) = A(t)x(t) + B(t)u(t),$$

where $x(\cdot) \in \mathbb{R}^n$ is a state vector, $u(\cdot) \in \mathbb{R}^m$ is a piecewise rd-continuous control vector function, $A(\cdot) \in \mathbb{R}^{n \times n}$ and $B(\cdot) \in \mathbb{R}^{n \times m}$ are rd-continuous matrices. (See [3], [1] and [2], for example). In order to investigate more general conditions for the functions $A(\cdot)$, $B(\cdot)$ and $u(\cdot)$, we consider a dynamic equation on times scales of the type

$$x^\Delta(t) = a(t)x(t) + b(t)u(t), \quad (1)$$

where $x(\cdot) \in X$ is a state vector, $a : [t_0, +\infty)_{\mathbb{T}} \rightarrow L(X)$ is Kurzweil-Henstock Δ -integrable (KH Δ -integrable), $u : [t_0, +\infty)_{\mathbb{T}} \rightarrow U$ and $b : [t_0, +\infty)_{\mathbb{T}} \rightarrow L(U, X)$ are such that $b(\cdot)u(\cdot)$ is KH Δ -integrable on $[t_0, +\infty)_{\mathbb{T}}$, and X, U, Y are Banach spaces. The goal of this work is to regard equation (1) as a generalized ordinary differential equation and to investigate necessary and sufficient conditions for the equation (1) to be controllable.

References

- [1] M. Bonher; A. Peterson, *Dynamic Equations on Times Scales: An Introduction with Applications*, Birkhäuser, Boston, 2001.
- [2] H. R. Henríquez; J. G. Mesquita, Self-Accessible states for linear system on time scales, *American Mathematical Society*, October, 2017.
- [3] K. Janglajew; E. Pawlussewics, Constrained local controllability of dinamic systems on time scales, *Advances id Difference Equations*, 2015.
- [4] V. Lupulescu; A. Younus, On controllability and observability for a class of linear impulsive dynamic systemas on time scales, *Mathematical and Computer Modelling*, 54, (2011), 130-1310.
- [5] A. Slavík, Dynamic equations on time scale and generalized ordinary differential equation, *J. Math. Anal. Appl.*, 385, (2012), 534-550.
- [6] E. Toon; F. A. Silva; M. Federson, Controllability and Observability for Linear Systems in Banach Spaces using Generalized Ordinary Differential Equations, pre-print.
- [7] E. Toon; F. A. Silva; M. Federson, Controllability for dynamic systems on time scales via Kurzweil-equations, pre-print.

Optimal Control on Time Scales: Necessary Optimality Conditions

Valeriano Antunes de Oliveira

Partially supported by grants 2013/07375-0 and 2016/03540-4 from FAPESP,
and by grants 457785/2014-4 and 310955/2015-7 from CNPq

Geraldo Nunes Silva

Partially supported by grant 2013/07375-0 from FAPESP through
Center for Mathematical Sciences Applied to Industry CEMEAI-CEPID

e-mail: geraldo.silva@unesp.br

Department of Applied Mathematics
UNESP – State University of São Paulo
São José do Rio Preto, São Paulo, Brasil

Jaqueline Godoy Mesquita

Department of Mathematics
UnB – Universidade de Brasília
Brasília, Distrito Federal, Brasil

Abstract

We consider smooth optimal control problems on time scales and provide necessary optimality conditions in the form of the Maximum Principle. The result presented here are under the convexity assumption on the velocity set, besides the usual assumptions. Although the convexity is an assumption which is little bit restrictive we expect to remove it in forthcoming works.

Periodic functions on isolated Time Scales

Bohner, Martin

e-mail: bohner@mst.edu

Department of Mathematics & Statistics
Missouri University of Science and Technology
Rolla, MO, USA

Mesquita, Jaqueline G.

e-mail: jgmesquita@unb.br

Departamento de Matemática
Universidade de Brasília, Campus Universitário Darcy Ribeiro
Asa Norte, Brasília-DF, Brazil

Streipert, Sabrina

e-mail: s.streipert@uq.edu.au

Centre for Applications in Natural Resource Mathematics
School of Mathematics and Physics, University of Queensland
St Lucia, QLD, Australia

Abstract

In this work, we formulate the definition of periodicity for isolated time scales. This provides the base for future research regarding periodicity on time scales with a positive graininess. The introduced definition is consistent with the known formulations in the discrete and quantum calculus setting. Using the definition of periodicity, we discuss the existence and uniqueness of ω -periodic solutions to a family of linear dynamic equations on isolated time scales.

References

- [1] M. Bohner and R. Chieochan. Floquet theory for q -difference equations. *Sarajevo J. Math.*, **8**(21)(2):355–366, 2012.
- [2] M. Bohner and J. G. Mesquita. Massera’s theorem in quantum calculus. *Proc. Amer. Math. Soc.*, 2018. To appear.
- [3] M. Bohner and A. Peterson. *Dynamic equations on time scales: An introduction with applications*. Birkhäuser Boston, Inc., Boston, MA, 2003.
- [4] M. Bohner and S. Streipert. Optimal harvesting policy for the Beverton-Holt quantum difference model. *Math. Morav.*, **20**(2):39-57, 2016.

Lambert W for time scales

Tom Cuchta

Department of Computer Science and Math
Fairmont State University
Fairmont, WV, USA

Abstract

The classic Lambert W function is a well-known special function that inverts the map $z \mapsto ze^z$. We investigate a time scales calculus generalization of the Lambert W function that inverts the map $z \mapsto ze_z(t, t_0)$, where $e_z(t, t_0)$ denotes the exponential function on a time scale. Special emphasis will be given to the time scale $\mathbb{T} = \mathbb{Z}$ (i.e. difference equations).

References

- [1] M. Bohner and Allan Peterson. *Dynamic equations on time scales*. Birkhäuser Boston Inc., Boston, MA, 2001. An introduction with applications.
- [2] R.M. Corless, G.H. Gonnet, D.E.G. Hare, D.J. Jeffrey, and D.E. Knuth. *On the Lambert W function*. *Adv. Comput. Math.*, 5(4):329–359, 1996.
- [3] Robert M. Corless, David J. Jeffrey, and Donald E. Knuth. *A sequence of series for the Lambert W function*. In *Proceedings of the 1997 international symposium on symbolic and algebraic computation, ISSAC '97, Maui, HI, USA, July 21–23, 1997*, pages 197–204. New York, NY: ACM Press, 1997.
- [4] D.J. Jeffrey, D.E.G. Hare, and Robert M. Corless. *Unwinding the branches of the Lambert W function*. *Math. Sci.*, 21(1):1–7, 1996.
- [5] István Mező and Árpád Baricz. *On the generalization of the Lambert W function*. *Trans. Am. Math. Soc.*, 369(11):7917–7934, 2017.
- [6] Seán M. Stewart. *On certain inequalities involving the Lambert W function*. *JIPAM, J. Inequal. Pure Appl. Math.*, 10(4):4, 2009.

Ordinary/Functional Differential Equations

Organizer: Pierluigi Benevieri (USP)

On systems of fractional differential equations with a general derivative

Agnieszka B. Malinowska

Supported by Bialystok University of Technology grant S/WI/1/2016

Faculty of Computer Science

Bialystok University of Technology

Bialystok, Poland

Abstract

Fractional Differential Equations (FDE) are a generalization of ordinary differential equations, where integer-order derivatives are replaced by fractional derivatives. Over the last decades, those equations have attracted a lot of attention of researchers from different areas, since fractional derivatives provide an excellent tool for the description of memory and hereditary properties of various materials and processes. Several types of fractional derivatives have been suggested in order to describe more accurately the real world phenomena, each ones with their own advantages and disadvantages. A more general unifying perspective to the subject was proposed in [2], by considering fractional operators depending on general kernels. In this work, we follow the special case of this approach that was developed in [1]. This type of differentiation depends on a kernel ϕ , and for some particular choices of ϕ , we obtain the well known fractional derivatives like the Caputo or the Caputo–Hadamard fractional derivatives.

We present the results on the existence and uniqueness of solutions to systems of FDEs with a derivative that depends on a kernel ϕ . Then we address the leader-follower consensus tracking problem for multi-agent systems. The consensus algorithm is proposed and its convergence is proved based on the stability results for linear systems of FDEs with a general form of fractional derivative. The effectiveness of the theoretical results is demonstrated through numerical simulations.

The presented results are based on the joint works with R. Almeida and T. Odziejewicz.

References

- [1] R. Almeida. *A Caputo fractional derivative of a function with respect to another function*. Commun. Nonlinear Sci. Numer. Simul., **44**, 460–481 (2017).
- [2] A. B. Malinowska, T. Odziejewicz and D. F. M. Torres. *Advanced methods in the fractional calculus of variations*. Springer Briefs in Applied Sciences and Technology, Springer, Cham, 2015.

Beyond integrability: real life eigenvalue algorithms

Carlos Tomei

Partially supported by CNPq and FAPERJ,

e-mail: carlos.tomei@gmail.com

Departamento de Matemática

PUC-Rio

Rio de Janeiro, Brazil

Abstract

In the eighties, the celebrated QR method, used to compute eigenvalues of matrices, has been shown to be interpolated by a flow in the integrable hierarchy of the Toda lattice. Shift techniques improve performance dramatically. We consider the dynamics of these algorithms. Joint work with Nicolau Saldanha (PUC-RIO) and Ricardo Leite (UFES).

References

- [1] P. Deift, T. Nanda, and C. Tomei, *Ordinary Differential Equations and the Symmetric Eigenvalue Problem*, SIAM J. Numer. Anal., 20(1), 1-22, 1983.

Elliptic equations and systems with critical Trudinger-Moser nonlinearities

J.M. do Ó
Departament of Mathematics
Brasília University
Brasília, DF, Brazil

Abstract

Existence of solutions is established for a class of systems of coupled equations via variational methods. Typical features of this class of problems are the lack of compactness because the unboundedness of the domain and critical growth. Moreover, the associated functional is strongly indefinite. The main tool used is the Trudinger-Moser inequality combined with a linking theorem.

References

- [1] D. de Figueiredo, J.M. do Ó, B. Ruf, *Elliptic equations and systems with critical Trudinger-Moser nonlinearities*, Discrete Contin. Dyn. Syst. **30** (2011), 455–476.
- [2] M. de Souza, J.M. do Ó, *Hamiltonian elliptic systems in \mathbb{R}^2 with subcritical and critical exponential growth*, Ann. Mat. Pura Appl. **195** (2016), 935–956.
- [3] J.M. do Ó, F. Sani, J. Zhang, *Stationary nonlinear Schrödinger equations in \mathbb{R}^2 with potentials vanishing at infinity*, Ann. Mat. Pura Appl. **196** (2017), 363–393.

Bifurcation theory for generalized ODE's

Maria Carolina Mesquita, Márcia Federson, Karina Schiabel

Partially supported by CAPES

e-mail: mc12stefani@hotmail.com

Department of Mathematics

Federal University of São Carlos

São Carlos, Brazil

Abstract

We establish conditions on the existence of bifurcation points of solutions of generalized ordinary differential equations via coincidence degree theory. We also present applications to ordinary differential equations.

References

- [1] Š. Schwabik *Abstract Perron-Stieltjes integrals*. Math. Bohem., **121**, 1996.
- [2] J. Kurzweil *Generalized ordinary differential equations*. Czechoslovak Math, 1958.
- [3] Š. Schwabik *Linear Stieltjes integral equations in Banach spaces*. Math. Bohem., **124**, 1999.
- [4] R. E. Gaines ; J. Mawhin *Coincidence Degree and Nonlinear Differential Equations* Lecture Notes in Math., **568**, Springer-Verlag, 1977.

On oscillation of Functional Differential Equations with delay with values in \mathbb{R}^n

Marielle Aparecida Silva
Partially supported by CAPES
e-mail: marielle@usp.br
ICMC
University of São Paulo
São Carlos, Brazil

Abstract

Consider the mesuare differential equation with impulses and delay

$$\begin{cases} (Dy_1(t), \dots, Dy_n(t)) = -(p_1(t)y_1(t - \tau_1), \dots, p_n(t)y_n(t - \tau_n))Dg \\ y_j(t_k^+) - y_j(t_k) = b_k y_j(t_k), \quad k \in \mathbb{N}, \quad j = 1, \dots, n, \end{cases} \quad (2)$$

where $\tau_j > 0$ is a constant, $p_j : [t_0, \infty) \rightarrow \mathbb{R}$ are functions, $j = 1, \dots, n$, $g : [t_0, \infty) \rightarrow \mathbb{R}$ is a regulated function which is left-continuous and continuous at the points of impulses t_k and Dy_j and Dg stand for the distributional derivatives of the functions y_j and g in the sense of distributions of L. Schwartz and, moreover,

- $t_0 < t_1 < \dots < t_k < \dots$ are fixed points and $\lim_{k \rightarrow \infty} t_k = \infty$;
- for $k \in \mathbb{N}$, $b_k \in (-\infty, -1) \cup (-1, \infty)$ are constants;
- for each compact subset $[a, b]$ of $[t_0, \infty)$ and $j = 1, \dots, n$, the Perron-Stieltjes integral $\int_a^b p_j(s) dg(s)$ exists.

The goal of this work is to present new criteria for the existence of oscillatory and nonoscillatory solutions of (1), that is, we propose oscillation and nonoscillation criteria for functional differential equations which the functions involved take values in \mathbb{R}^n .

References

- [1] Silva, M. Ap.; Federson, M.; Gadotti, M. C., *Oscillation and nonoscillation criteria for impulsive delay differential equations with Perron integrable righthand sides*. Submitted

Uniform persistence and periodic solutions for some systems of delay differential equations

Pablo Amster

Partially supported by CONICET, PIP 11220130100006CO

e-mail: pamster@dm.uba.ar

Departamento de Matemática - FCEyN

Universidad de Buenos Aires & IMAS-CONICET

Buenos Aires, Argentina

Abstract

Simple criteria for persistence in some systems of delay differential equations shall be presented. Inspired by a classical population model, we shall introduce sufficient conditions for weak or strong persistence and uniform persistence. As a consequence, we shall deduce the existence of positive periodic solutions. Moreover, we shall prove, in some cases, that the conditions are also necessary; in particular, accurate assumptions will imply that 0 is a global attractor for all the positive solutions.

References

- [1] P. Amster, A. Déboli, *Necessary and Sufficient Conditions for the Existence of Periodic Solutions of a Nicholson Type Delay System*. *Differ. Equ. Dyn. Syst.* (2016), <https://doi.org/10.1007/s12591-016-0285-y>
- [2] L. Berezansky, E. Braverman, L. Idels, *Nicholson's blowflies differential equation revisited: main results and open problems*. *Appl. Math. Model.* **34**, (2010) 1405–1417.
- [3] J. So, J. S. Yu, *Global attractivity and uniform persistence in Nicholson's blowflies*, *Diff. Eqns. Dynam. Syst.* **2** (1) (1994) 11–18

Global persistence of nonlinear operators in Hilbert spaces

Pierluigi Benevieri

e-mail: pluigi@ime.usp.br

Instituto de Matemática e Estatística

Universidade de São Paulo – USP, Brasil

Abstract

We study the nonlinear eigenvalue problem $Lx + \varepsilon N(x) = \lambda x$ in a finite-dimensional real Hilbert space H , where ε, λ are real parameters, $L: H \rightarrow H$ is linear, and $N: S \rightarrow H$ is a continuous map, S being the unit sphere in H . We prove a global continuation property of the set of the *solution triples* $(x, \varepsilon, \lambda) \in S \times \mathbb{R} \times \mathbb{R}$ of this problem. Namely, under the assumption that l_* is a simple eigenvalue of L with eigenvector $x_* \in S$, we show that, in the set of all the solution triples, the connected component containing $(x_*, 0, \lambda_*)$ is either unbounded or meets a solution triple $(x^*, 0, l^*)$ with $x^* \neq x_*$.

Our results are related to some papers by R. Chiappinelli, in which he studied a “local” persistence property of eigenvalues and eigenvectors of self-adjoint operators in real Hilbert spaces. This is a joint work with A. Calamai, M. Furi, M.P. Pera

References

- [1] Benevieri P. - Calamai A. - Furi M. - Pera M.P., *On a result of Chiappinelli about the persistence of the eigenvalues of a perturbed self-adjoint operator*, preprint.
- [2] Benevieri P. - Calamai A. - Furi M. - Pera M.P., *On the persistence of the eigenvalues of a perturbed Fredholm operator of index zero under nonsmooth perturbations*, *Z. Anal. Anwend.* **36** (2017), no. 1, 99–128.
- [3] Benevieri P. - Calamai A. - Furi M. - Pera M.P., *Global continuation of the eigenvalues of a perturbed linear operator*, *Annali di Mat. Pura ed Appl.* DOI 10.1007/s10231-017-0717-5.
- [4] Chiappinelli R., *Isolated Connected Eigenvalues in Nonlinear Spectral Theory*, *Nonlinear Funct. Anal. Appl.* **8** (2003), no. 4, 557–579.
- [5] Chiappinelli R. - Furi M. - Pera M.P., *Normalized Eigenvectors of a Perturbed Linear Operator via General Bifurcation*, *Glasg. Math. J.* **50** (2008), no. 2, 303–318.
- [6] Chiappinelli R. - Furi M. - Pera M.P., *Topological persistence of the normalized eigenvectors of a perturbed self-adjoint operator*, *Appl. Math. Lett.* **23** (2010), no. 2, 193–197.
- [7] Chiappinelli R. - Furi M. - Pera M.P., *Persistence of the normalized eigenvectors of a perturbed operator in the variational case*, *Glasg. Math. J.* **55** (2013), no. 3, 629–638.
- [8] Chiappinelli R. - Furi M. - Pera M.P., *Topological Persistence of the Unit Eigenvectors of a Perturbed Fredholm Operator of Index Zero*, *Z. Anal. Anwend.* **33** (2014), no. 3, 347–367.

Optimality Conditions for Optimal Control Problems with Mixed Constraints

Valeriano Antunes de Oliveira

Partially supported by grants 2013/07375-0 and 2016/03540-4 from FAPESP,
and by grants 457785/2014-4 and 310955/2015-7, from CNPq,

e-mail: valeriano.oliveira@unesp.br

Geraldo Nunes Silva

Department of Applied Mathematics
UNESP – State University of São Paulo
São José do Rio Preto, São Paulo, Brasil

Jamielli Tomaz Pereira, Roberto Andreani

Department of Applied Mathematics
UNICAMP – State University of Campinas
Campinas, São Paulo, Brazil

Abstract

Necessary and sufficient optimality conditions are given for optimal control problems with mixed state-control constraints. The necessary conditions are given in the form of a weak maximum principle and are obtained under a constant rank type regularity condition. Such necessary conditions are shown to be also sufficient for optimality under generalized convexity assumptions. The concept of generalized convexity is the most general one.

Problem statement and optimality conditions

The optimal control problem with mixed constraints this work is concerned with is posed as follows:

$$\begin{aligned}
 & \text{minimize} && l(x(0), x(1)) \\
 & \text{subject to} && \dot{x}(t) = f(t, x(t), u(t), v(t)) \text{ a.e. in } [0, 1], \\
 & && h(t, x(t), u(t), v(t)) = 0 \text{ a.e. in } [0, 1], \\
 & && g(t, x(t), u(t), v(t)) \leq 0 \text{ a.e. in } [0, 1], \\
 & && v(t) \in V(t) \text{ a.e. in } [0, 1], \\
 & && (x(0), x(1)) \in C,
 \end{aligned} \tag{P}$$

where $l : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$, $(f, g, h) : [0, 1] \times \mathbb{R}^n \times \mathbb{R}^{k_u} \times \mathbb{R}^{k_v} \rightarrow \mathbb{R}^n \times \mathbb{R}^{m_g} \times \mathbb{R}^{m_h}$ are given functions, $V(t) \subset \mathbb{R}^{k_v}$ for all $t \in [0, 1]$ and $C \subset \mathbb{R}^n \times \mathbb{R}^n$.

Let us denote the unmaximized Hamiltonian function as

$$H(t, x, p, q, r, u, v) := p \cdot f(t, x, u, v) + q \cdot h(t, x, u, v) + r \cdot g(t, x, u, v).$$

The weak maximum principle is said to be satisfied at a feasible process $(\bar{x}, \bar{u}, \bar{v})$ if there exist $p \in W^{1,1}([0, 1]; \mathbb{R}^n)$, $q \in L^1([0, 1]; \mathbb{R}^{m_h})$, $r \in L^1([0, 1]; \mathbb{R}^{m_g})$, $\zeta \in L^1([0, 1]; \mathbb{R}^{k_v})$, and $\lambda \geq 0$ such that

- (i) $\|p\|_\infty + \lambda \neq 0$;
- (ii) $(-\dot{p}(t), 0, \zeta(t)) = \nabla_{x,u,v} H(t, \bar{x}(t), p(t), q(t), r(t), \bar{u}(t), \bar{v}(t))$ a.e. in $[0, 1]$;
- (iii) $\zeta(t) \in \text{co}N_{V(t)}(\bar{v}(t))$ a.e. in $[0, 1]$;
- (iv) $r(t) \cdot g(t, \bar{x}(t), \bar{u}(t), \bar{v}(t)) = 0$ and $r(t) \leq 0$ a.e. in $[0, 1]$;

$$(v) \quad (p(0), -p(1)) \in N_C(\bar{x}(0), \bar{x}(1)) + \lambda \nabla l(\bar{x}(0), \bar{x}(1)).$$

Furthermore, for some integrable function K_m , $|(q(t), r(t))| \leq K_m(t)|p(t)|$ a.e. in $[0, 1]$.

In this work, necessary optimality conditions are given for (P) in the form of the weak maximum principle stated above, where a regularity condition of constant rank type is imposed on the mixed constraints. This regularity condition is weaker than full rank assumptions found in the literature (see de Pinho [3] and references therein), and is an alternative regularity condition regarding Mangasarian-Fromovitz constraint qualification (see de Pinho and Rosenblueth [4]). Sufficient optimality conditions are obtained in terms of the so called maximum-principle-pseudo-invexity (MP-pseudo-invexity) (see de Oliveira and Silva [1, 2]). As expected, MP-pseudo-invex problems subsume convex problems. Moreover, it is shown that when all extremal control processes are optimal, the problem is, necessarily, MP-pseudo-invex. This means that the class of MP-pseudo-invex problems is the largest class of optimal control problems in which the optimality conditions of the maximum principle are both necessary and sufficient.

References

- [1] V. A. de Oliveira and G. N. Silva. *New optimality conditions for nonsmooth control problems*. J. Glob. Optim., **57**, 1465–1484, 2013.
- [2] V. A. de Oliveira and G. N. Silva. *On sufficient optimality conditions for multiobjective control problems*. J. Glob. Optim., **64**, 721–744, 2016.
- [3] M. R. de Pinho. *Mixed constrained control problems*. J. Math. Anal. Appl., **278**, 293–307, 2003.
- [4] M. R. de Pinho and J. F. Rosenblueth. *Necessary conditions for constrained problems under Mangasarian-Fromovitz conditions*. Siam J. Control Optim., **47**, 535–552, 2008.

Nonlinear Dynamical Systems

Organizer: Juliana Pimentel (UFABC)

The Variational Principle for Locally Compact Spaces

André Caldas de Souza
e-mail: andrecaldas@unb.br
Department of Mathematics
University of Brasília
Brasília, Brazil

Abstract

We have extended the *variational principle* for entropies to continuous transformations $T : X \rightarrow X$ defined over locally compact separable metrizable spaces. To each T -invariant Radon probability measure μ , there is an associated *Kolmogorov-Sinai* entropy, $h_\mu(T)$. The *Bowen entropy*, $h_d(T)$, is defined for each metric d compatible with the topology of X . And when X is compact, Adler, Konheim e McAndrew have defined the *topological entropy*, $h(T)$, in terms of open coverings. To deliver a theory similar to the compact case (Theorem 8.6 in [2]), we have extended in [1] the concept of *topological entropy* using open coverings such that at least one of its members has compact complement. Using this new definition, we have shown that

$$\sup_{\mu} h_{\mu}(T) = h(T) = \min_d h_d(T),$$

where the minimum of $h_d(T)$ is attained when d is the restriction to X of any metric defined over the *one point compactification* of X .

References

- [1] A. Caldas & M. Patro. *Entropy and its variational principle for locally compact metrizable systems*. Ergodic Theory and Dynamical Systems, **38**(2):540–565, 2018.
- [2] P. Walters. *An Introduction to Ergodic Theory*. Springer-Verlag, 2000.

Impulsive non-autonomous dynamical systems

Everaldo de Mello Bonotto

Partially supported by FAPESP grant 2014/25970-5 and CNPq grant 307317/2013-7

e-mail: ebonotto@icmc.usp.br

Department of Applied Mathematics and Statistics

University of São Paulo

São Carlos, Brazil

Abstract

In this talk, we present the theory of “impulsive non-autonomous dynamical systems”. We establish conditions to ensure the existence of an impulsive cocycle attractor and we present some examples.

References

- [1] E. M. Bonotto, M. C. Bortolan, T. Caraballo and R. Collegari, *Impulsive non-autonomous dynamical systems and impulsive cocycle attractors*, Math. Meth. Appl. Sci., **40**, 1095-1113, 2017.

Construction of a global solution for a process

Jacson Simsen

Partially supported by FAPEMIG (Brazil) - processes PPM 00329-16 and CEX-APQ-00814-16.

e-mail: jacson@unifei.edu.br

Institute of Mathematics and Computation

Federal University of Itajubá (UNIFEI)

Itajubá, Brazil.

Abstract

Construction of bounded complete trajectories for semigroups (see [1, 2]) and bounded complete orbits for multivalued semiflows or generalized semiflows (see [3]) is an old subject of extreme importance for characterization of attractors . I will present in this talk a technique developed in [4] to construct a bounded global solution for a process in order to obtain upper semicontinuity of a family of pullback attractors.

References

- [1] J.K. Hale, *Asymptotic behavior of dissipative systems*. Mathematical Surveys and Monographs, 25. American Mathematical Society, Providence, RI, 1988.
- [2] O. Ladyzhenskaya, *Attractors for semigroup and evolution equations* . Cambridge University Press, 1991.
- [3] J.M. Ball, Continuity properties and global attractors of generalized semiflows and the Navier-Stokes equations. *J. Nonlinear Sci.* 7 (1997), no. 5, 475–502.
- [4] A.C. Fernandes, M.C. Gonçalves and J. Simsen, Non-autonomous Reaction-Diffusion equations with variable exponents and large diffusion. To appear at *Discrete and Continuous Dynamical Systems - B*, 2018.

Combining random searches, individual's memory, and population dynamics: analysis of population redistribution patterns

Juliana M. Berbert

Partially supported by by FAPESP (#2014/17833-8)

e-mail: juliana.berbert@ufabc.edu.br

Karen A. Oliveira

Partially supported by CNPq

e-mail: karen.amaral.oliveira@gmail.com

Centro de Matemática, Computação e Cognição
Universidade Federal do ABC, Santo André, SP, Brazil.

Abstract

The reaction-diffusion equation is one of the possible ways for modeling animal movement, where the reactive part stands for the population growth and the diffusive part for random dispersal of the population. Nevertheless, there are some mechanisms that affect the movement, such as resource depletion or individual's spatial memory of recently depleted patches, which results in a bias for one direction of dispersal and can be modeled by an advective term on a advection-reaction-diffusion equation. Therefore, our model is composed of a coupled nonlinear partial differential equation system with one equation for the population dynamics and other for the memory density distribution. For the population growth, we use either the exponential or logistic growth function. Analytic approach, shows that for the exponential and logistic growth the traveling wave speeds are the same with or without memory dynamics. From numerical analysis, we explore how the population redistribution is affected by different values of the parameters: memory, growth rate, and carrying capacity. Combining these parameters results on a redistribution pattern of the population associated to either normal or anomalous diffusion: subdiffusion and superdiffusion.

References

- [1] J.M. Berbert, and M.A. Lewis. *Superdiffusivity due to resource depletion in random searches*. Ecological Complexity, **33**, p. 41-48, 2018.

The minimal Morse components of translations in flag manifolds are normally hyperbolic

Lucas Seco

e-mail: `lsec@unb.br`

Department of Mathematics

University of Brasília

Brasília, Brasil

Mauro Patrão

Department of Mathematics

University of Brasília

Brasília, Brasil

Abstract

In the projective space, consider the iteration of an invertible matrix: are the minimal Morse components normally hyperbolic? An affirmative answer was only known in the case where the matrix is diagonalizable. Using Lie theory, we prove this to be true in the rather more general context of an arbitrary element of a semisimple Lie group acting on one of its flag varieties: the so-called translations in flag varieties. This context encompasses the iteration of invertible matrices not necessarily diagonalizable in the real or complex projective space, the classic flag varieties of real or complex nested spaces, and also the symplectic grassmanians. Without using the tools from Lie Theory we do not know a solution for this problem even in the simplest case of the projective space. In this lecture we will sketch how we solve this question for the projective space, presenting the necessary rudiments from Lie theory.

References

- [1] M. Patrão, L. Seco: *The minimal Morse components of translations on flag manifolds are normally hyperbolic*, submitted.
- [2] T. Ferraiol, M. Patrão, L. Seco: *Jordan decomposition and dynamics on flag manifolds*, Discrete Contin. Dynam. Systems A, **26**(3) (2010).
- [3] C. Pugh and M. Shub: *Linearization of normally hyperbolic diffeomorphisms and flows*. Invent. Math. **10** (1970), 187-198.
- [4] M. Shub and A. Vasquez: *Some linearly induced Morse-Smale systems, the QR algorithm and the Toda lattice*, Contemp. Math., **64** (1987), 181-194.
- [5] G. Ammar and C. Martin: *The Geometry of Matrix Eigenvalues Methods*, Acta Appl. Math. **5** (1986), 239-278.

The Topological Entropy of Endomorphisms of Lie Groups

Mauro Patrão

e-mail: mpatrao@mat.unb.br

Department of Mathematics

University of Brasília, Brazil.

Abstract

In this talk, we determine the topological entropy $h(\phi)$ of a continuous endomorphism ϕ of a Lie group G . This computation is a classical topic in ergodic theory which seemed to have long been solved. But, when G is noncompact, the well known Bowen's formula for the entropy $h_d(\phi)$ associated to a left invariant distance d just provides an upper bound to $h(\phi)$, which is characterized by the so called variational principle. We prove that

$$h(\phi) = h\left(\phi|_{T(G_\phi)}\right)$$

where G_ϕ is the maximal connected subgroup of G such that $\phi(G_\phi) = G_\phi$, and $T(G_\phi)$ is the maximal torus in the center of G_ϕ . This result shows that the computation of the topological entropy of a continuous endomorphism of a Lie group reduces to the classical formula for the topological entropy of a continuous endomorphism of a torus. Our approach explores the relation between null topological entropy and the nonexistence of Li-Yorke pairs and also relies strongly on the structure theory of Lie groups.

References

- [1] R. Bowen: *Entropy for group endomorphisms and homogeneous spaces*. Trans. Americ. Math Soc. **153** (1971), 401-414.
- [2] A. Caldas and M. Patrão: *Entropy of Endomorphisms of Lie Groups*. Discrete and Continuous Dynamical Systems **33** (2013), 1351-1363.
- [3] A. Caldas and M. Patrão: *Entropy and Its Variational Principle for Locally Compact Metrizable Systems*. Ergodic Theory and Dynamical Systems (2016), 1-26.
- [4] M. Patrão: *Entropy and its variational principle for non-compact metric spaces*. Ergodic Theory and Dynamical Systems **30** (2010), 1529-1542.

A nonlocal approach to spatial spread in thin structures

Marcone C. Pereira

Partially supported by CNPq 303253/2017-7, FAPESP 2017/02630-2

e-mail: marcone@ime.usp.br

Department of Applied Mathematics

University of São Paulo

São Paulo, Brazil

Julio D. Rossi

Partially supported by MINCYT grant MTM2016-68210 (Spain)

e-mail: jrossi@dm.uba.ar

Department of Mathematics

University of Buenos Aires

Buenos Aires, Argentina

Abstract

In this talk we discuss an approach to considerer spatial spread in N -dimensional thin structures. We introduce equations with nonlocal dispersal and defined in tight domains contrasting it with its corresponding local diffusion equation with Neumann and Dirichlet boundary conditions. Here the thin structure effect is modeled by an ϵ -parameter family of open sets which squeezes to a lower dimension open set as $\epsilon \rightarrow 0$. The asymptotic behavior of the solutions is analyzed and the results are compared with classical situations to elliptic equations in thin domains.

References

- [1] M. C. Pereira and J. D. Rossi. *Nonlocal problems in thin domains*; J. of Diff. Eq. 263 (2017) 1725-1754.
- [2] M. C. Pereira and J. D. Rossi. *Nonlocal evolution problems in thin domains*; Applicable Analysis DOI 10.1080/00036811.2017.1350850.

Unbounded Sturm attractors for quasilinearparabolic equations

Phillipo Lappicy
Departament of Mathematics
University of São Paulo
São Carlos, Brazil

Abstract

The goal of this paper is to construct explicitly the global attractors of quasi-linear parabolic equations when solution can also grow up, and hence the attractor is unbounded. In particular, we construct heteroclinic connections between bounded and/or unbounded hyperbolic equilibria. This is a joint work with Juliana Pimentel.

Rayleigh–Bénard problem for micropolar fluids

Piotr Kalita

e-mail: piotr.kalita@ii.uj.edu.pl

Faculty of Mathematics and Computer Science

Jagiellonian University

Kraków, Poland

Abstract

The talk will be devoted to the Rayleigh–Bénard convection of a micropolar fluid. Convective flows in such fluids are governed by the following system of equations

$$\begin{aligned}\frac{1}{\text{Pr}}(u_t + (u \cdot \nabla)u) - (1 + K)\Delta u + \frac{1}{\text{Pr}}\nabla p &= 2K\text{rot } \omega + e_3\text{Ra}T \\ \text{div } u &= 0 \\ \frac{M}{\text{Pr}}(\omega_t + (u \cdot \nabla)\omega) - L\Delta\omega - G\nabla\text{div } \omega + 4K\omega &= 2K\text{rot } u \\ T_t + u \cdot \nabla T - \Delta T &= 0,\end{aligned}$$

where, in contrast to Newtonian fluids, in order to close the system, one needs to consider the additional equation for the angular momentum ω . We will present the results on the existence and dimension of global attractor and compare them with the corresponding results for Newtonian fluids. In three-dimensional case we will show that the global attractor exists for large Prandtl number Pr . We will also show the estimates for the Nusselt number obtained by the background flow method, for finite and infinite Prandtl number, and compare them with the known estimates for the Newtonian fluids. The estimates for micropolar fluids turn out to be sharper than the corresponding ones for Newtonian fluids, which reflects the fact that additional dissipation effects related with the rotational friction between fluid particles are present in the model.

Robustness with respect to exponents for non-autonomous reaction diffusion equations

Rodrigo A. Samproгна

Partially supported by CAPES process 1728825

e-mail: rodrigo.samproгна@unifal-mg.edu.br

Instituto de Ciência e Tecnologia

Universidade Federal de Alfenas campus Poços de Caldas

Poços de Caldas, Brazil

Jacson Simsen

Partially supported by FAPEMIG processes PPM 00329-16

e-mail: jacson@unifei.edu.br

Instituto de Matemática e Computação

Universidade Federal de Itajubá

Itajubá, Brazil

Abstract

In this work we establish upper semicontinuity of pullback attractors for a non-autonomous evolution equation of the form

$$\begin{cases} \frac{\partial u_\lambda}{\partial t}(t) - \operatorname{div} (D(t)|\nabla u_\lambda(t)|^{p_\lambda(x)-2}\nabla u_\lambda(t)) + |u_\lambda(t)|^{p_\lambda(x)-2}u_\lambda(t) = B(t, u_\lambda(t)), \\ u_\lambda(\tau) = u_{0\lambda} \end{cases} \quad (P_\lambda)$$

with a homogeneous Neumann boundary condition, for $(t, x) \in (\tau, +\infty) \times \Omega$ where Ω is a bounded smooth domain in \mathbb{R}^N for some $N \geq 1$ and the initial condition $u_\lambda(\tau) \in H := L^2(\Omega)$. The exponents functions $p_\lambda \in C(\overline{\Omega}, \mathbb{R})$ satisfy some control assumptions and $p_\lambda \rightarrow p$ in $L^\infty(\Omega)$ for some p such that $p(\cdot) \in C(\overline{\Omega}, \mathbb{R})$.

The terms B and D are assumed to satisfy:

Assumption B The mapping $B : [\tau, T] \times H \rightarrow H$ is such that

(B1) there exist $L \geq 0$ such that

$$\|B(t, x_1) - B(t, x_2)\|_H \leq L\|x_1 - x_2\|_H$$

for all $t \in [\tau, T]$ and $x_1, x_2 \in H$;

(B2) for all $x \in H$ the mapping $t \rightarrow B(t, x)$ belongs to $L^2(\tau, T; H)$;

(B3) the function $t \rightarrow \|B(t, 0)\|_H$ is nondecreasing, absolutely continuous and bounded on compact subsets of \mathbb{R} .

Assumption D $D : [\tau, T] \times \Omega \rightarrow \mathbb{R}$ is a function in $L^\infty([\tau, T] \times \Omega)$ such that

(D1) there are positive constants, β and \mathcal{M} such that $0 < \beta \leq D(t, x) \leq \mathcal{M}$ for almost all $(t, x) \in [\tau, T] \times \Omega$;

(D2) $D(t, x) \geq D(s, x)$ for each $x \in \Omega$ and $t \leq s$ in $[\tau, T]$.

The authors in [1] also considered the non-autonomous problem and proved the robustness with respect to the diffusion coefficient whereas in this work we study the robustness with respect to the exponents. The results of this research can be found in detail in [2].

References

- [1] P. E. Kloeden, J. Simsen. *Pullback attractors for non-autonomous evolution equations with spatially variable exponents*. Commun. Pure Appl. Anal., **13**, no. 6, 2014.
- [2] R. A. Samprogna, J. Simsen. *Robustness with respect to exponents for non-autonomous reaction diffusion equations*. Electron. J. Qual. Theory Differ. Equ., **no. 11**, 2018.

Intermingled basins in coupled Lorenz systems

Sabrina Camargo

e-mail: sabrina.camargo@ufabc.edu.br

Centro de Matemática, Computação e Cognição

Universidade Federal do ABC

Santo André, Brasil

Ricardo L. Viana

Departamento de Física

UFPR

Curitiba, Brasil

Celia Anteneodo

Departamento de Física

PUC-Rio

Rio de Janeiro, Brasil

Abstract

Riddled basins occur in nonlinear systems whose phase space symmetry allows an invariant subspace with a chaotic attractor. This invariant subspace can either attract or repel orbits. As a consequence, for every point belonging to the basin of attraction, there is another point, arbitrarily close, that does not belong to the basin of attraction. The presence of riddled basins is verified by analyzing the maximal transversal Lyapunov exponent and the maximal transversal finite time Lyapunov exponent.

In this work, we consider a system of two identical linearly coupled Lorenz oscillators presenting synchronization of chaotic motion for a specified range of the coupling strength. We verify the existence of global synchronization and antisynchronization attractors with intermingled basins of attraction such that the basin of one attractor is riddled with holes belonging to the basin of the other attractor and vice versa. We investigated this phenomenon by verifying the fulfillment of the mathematical requirements for intermingled basins and obtained scaling laws that characterize quantitatively the riddling of both basins in this system.

References

- [1] S. Camargo, R. L. Viana, C. Anteneodo *Intermingled basins in coupled Lorenz systems*. PRE, **85** 036207 (2012).

Functional Analysis

Organizer: Geraldo de Azevedo Botelho (UFU)

Duality in Spaces of Lorch Analytic Mappings

Luiza A. Moraes
e-mail: luiza@im.ufrj.br
Department of Mathematics
University Federal of Rio de Janeiro
Rio de Janeiro, Brazil

Alex F. Pereira
e-mail: alexpereira@id.uff.br
Department of Analysis
University Federal Fluminense
Niterói, Brazil

Abstract

Let E be a complex Banach algebra. In this talk we study the strong dual of the Fréchet space $\mathcal{H}_L(E)$ of the mappings $f : E \rightarrow E$ that are analytic in the sense of Lorch. In fact we are going to present two descriptions of the dual of the space $\mathcal{H}_L(E)$ and some recent results on topological properties of the strong dual of the space $\mathcal{H}_L(E)$.

References

- [1] J. M. Ansemil and S. Ponte. *An example of a quasi-normable Fréchet function space which is not a Schwartz space*. Advances in Functional Analysis, Holomorphy and Approximation Theory (Ed. S. Machado) Lecture Notes in Mathematics **843**, Springer Verlag, Berlin (1981) 1-8.
- [2] D. Garcia, M. L. Lourenço, L. A. Moraes and O. Paques. *The spectra of some algebras of analytic mappings*. Indag. Math. (N.S.) **10 (3)** (1999) 393-406.
- [3] A. Grothendieck. *Sur les espaces (F) et (DF)* . Summa Brasiliensis Math. **3** (1954) 57-122.
- [4] J. M. Isidro. *Topological duality on the functional space $(\mathcal{H}_b(U, F), \tau_b)$* . Proc. Roy. Irish Acad. **79A** (1979) 115-130.
- [5] L.A. Moraes and A.F. Pereira. *Duality in spaces of Lorch analytic mappings*. Quart. J. Math. **67** (2016) 431-438.

Twisted sums of c_0 and $C(K)$

Claudia Correa
Centro de Matemática, Computação e Cognição
Universidade Federal do ABC
Santo André, Brasil

Abstract

The purpose of this talk is to discuss a problem I have been working on for the last couple of years. This problem deals with the existence of nontrivial twisted sums of Banach spaces. It is an easy corollary of Sobczyk's Theorem that if X is a separable Banach space, then every twisted sum of c_0 and X is trivial. This naturally raises the question about the converse of this last implication, i.e., if X is a Banach space such that every twisted sum of c_0 and X is trivial, then X must be separable? This question is easily answered negatively. However it becomes quite interesting when we restrict ourselves to the subclass of Banach spaces of the form $C(K)$, i.e., the space of continuous real-valued functions defined on a compact Hausdorff space K , endowed with the supremum norm. Recall that a space $C(K)$ is separable if and only if K is metrizable. Therefore the question in this context can be rephrased as: Is there a nonmetrizable compact Hausdorff space K such that every twisted sum of c_0 and $C(K)$ is trivial? This question was proposed originally by Cabello, Castillo, Kalton and Yost in 2003 and it has not been solved yet. In this talk I will present the progress we have made towards the general solution of this problem as well as possible future developments.

References

- [1] F. Cabello, J. M. F. Castillo, N. J. Kalton & D. T. Yost, Twisted sums with $C(K)$ spaces, *Trans. Amer. Math. Soc.*, 355 (11), 2003, pgs. 4523—4541.
- [2] J. M. F. Castillo, Nonseparable $C(K)$ -spaces can be twisted when K is a finite height compact, *Topology Appl.*, 198, 2016, pgs. 107—116.
- [3] C. Correa & D. V. Tausk, Local extension property for finite height spaces, preprint: <https://arxiv.org/pdf/1801.08619.pdf>
- [4] C. Correa & D. V. Tausk, Nontrivial twisted sums of c_0 and $C(K)$, *J. Func. Anal.*, 270, 2016, pgs. 842—853.
- [5] W. Marciszewski & G. Plebanek, Extension operators and twisted sums of c_0 and $C(K)$ spaces, *J. Funct. Anal.*, 274 (5), 2018, pgs. 1491—1529.

On the spectrum of the algebra of bounded type holomorphic functions on Banach spaces

Daniela M. Vieira

Supported by FAPESP-Brazil, Proc. 2014/07373-0

e-mail: danim@ime.usp.br

Departament of Mathematics

University of So Paulo

So Paulo, Brazil

Abstract

We study the spectrum of the algebra of bounded type analytic functions on unit ball (or a complete Reinhardt domain in) in a Banach space E . We aim to describe $M_b(U)$ as analytic manifold. We show that whenever U is a complete Reinhardt domain in a reflexive space with 1-unconditional basis, each connected component of $M_b(U)$ is (identified with) a complete Reinhardt set. We also prove that, when U is the unit ball of ℓ_p , the connected components are identified with balls which, with the exception of the component formed by evaluations, have radius strictly smaller than 1. Moreover, there are connected components with arbitrary small radius. We also show that for other Banach sequence spaces, connected components do not necessarily identify with balls. Joint work with D. Carando and S. Muro, UBA, Argentina.

References

- [1] R. Aron, D. Carando, S. Lassalle, M. Maestre, *Cluster values of holomorphic functions of bounded type*, Trans. Amer. Math. Soc. **368** (2016) 2355-2369.
- [2] R. Aron, P. Galindo, D. Garcia, M. Maestre, *Regularity and algebras of analytic functions in infinite dimensions*, Trans. Amer. Math. Soc. **348** (1996) 543-559.
- [3] D. Carando, S. Muro, *Envelopes of holomorphy and extension of functions of bounded type*, Adv. Math. **229** (2012) 2098-2121.

Some stability properties of the polynomial daugavetian index

Elisa R. Santos

Partially supported by Fapemig Grant APQ-00522-14

e-mail: elisars@ufu.br

Faculty of Mathematics

Federal University of Uberlândia

Uberlândia, Brazil

Abstract

Let X be an infinite-dimensional complex Banach space and let P be a compact polynomial on X given by $P = P_0 + P_1 + \cdots + P_n$ where P_j is a j -homogeneous polynomial for $j = 0, \dots, n$. From [1, Proposition 3.4], we have that P_1 is a compact linear operator. And it follows from the Cauchy Inequality that

$$\|Id + P_1\| \leq \|Id + P\|.$$

Since P_1 is compact and X has infinity dimension, $\|Id + P_1\| \geq 1$ and consequently $\|Id + P\| \geq 1$. This allow us to define the *polynomial daugavetian index* of X as

$$\text{daug}_p(X) = \max \{m \geq 0 : \|Id + P\| \geq 1 + m\|P\| \text{ para todo } P \in \mathcal{P}_K(X)\},$$

generalizing the ideas of the daugavetian index defined by M. Martín [2]. Observe that $0 \leq \text{daug}_p(X) \leq 1$. When $\text{daug}_p(X) = 1$, the space X has the *polynomial Daugavet property*, that is, every weakly compact polynomial P on X satisfies the Daugavet equation:

$$\|Id + P\| = 1 + \|P\|.$$

The purpose of this talk is to present some stability properties of the daugavetian index for c_0 -, ℓ_∞ - and ℓ_1 -sums.

References

- [1] R. M. Aron, M. Schottenloher. *Compact Holomorphic Mappings on Banach Spaces and the Approximation Property*. J. Funct. Anal. **21** (1976), 7–30.
- [2] M. Martín. *The Daugavetian index of a Banach space*. Taiwanese J. Math. **7** (2003), 631–640.

Summability of multilinear operators: a unified theory and consequences

Gustavo Araújo

e-mail: gdsaraujo@gmail.com and gustavoaraujo@cct.uepb.edu.br

Departamento of Matemática
Universidade Estadual da Paraíba
Campina Grande, Brazil

Abstract

Results related to summability of multilinear operators date back, at least, to the 30's, when Littlewood proved his seminal $4/3$ inequality. Since then, several different related results and approaches have appeared, as the Hardy–Littlewood inequalities (Quarterly J. Math., 1934), that can be considered a keystones to the theory of summability of multilinear operators. In this talk we present a new unifying approach to a series of results on summability of multilinear forms, that include the aforementioned inequalities.

References

- [1] N. Albuquerque, G. Araújo, W. Cavalcante, T. Nogueira, D. Núñez-Alarcón, D. Pellegrino and P. Rueda, *On summability of multilinear operators and applications*, submitted, 2018.
- [2] G. Hardy and J.E. Littlewood, *Bilinear forms bounded in space $[p; q]$* , Quart. J. Math., **5**, 241-254, 1934.
- [3] R. Ryan, *Introduction to tensor products*, Springer Verlag, London, 2002.

Jorge Mujica's last theorem and applications

Geraldo Botelho

Partially supported by CNPq Grant 305958/2014-3 and Fapemig Grant PPM-00450-17

e-mail: botelho@ufu.br

Faculdade de Matemática

Universidade Federal de Uberlândia

Uberlândia, Brazil

Abstract

Let U be a balanced open subset of the complex Banach space E . Shortly before passing away quite prematurely, Jorge Mujica proved that the sequence $(Q(^nE))_{n=0}^{\infty}$, where $Q(^nE)$ is the space of complex-valued continuous n -homogeneous polynomials on E , is an \mathcal{S} -absolute Schauder decomposition for the predual $G_b(U)$ of the space $\mathcal{H}_b(U)$ of holomorphic functions of bounded type on U . In this talk we describe two applications of this result, concerning representations of linearizations of holomorphic functions of bounded type and closed ideals of linear operators between locally convex spaces. This is a joint work with Vinícius V. Fávoro and Jorge Mujica.

References

- [1] G. Botelho, V. V. Fávoro, J. Mujica. *Absolute Schauder decompositions and linearization of holomorphic mappings of bounded type*. Publ. Res. Inst. Math. Sci., to appear.

mid (q, p) -summable sequences

Jamilson R. Campos

e-mail: jamilson@dcx.ufpb.br or jamilsonrc@gmail.com

Departamento de Ciências Exatas

Universidade Federal da Paraíba

Rio Tinto, Brazil

Joedson Santos

e-mail: joedsonmat@gmail.com

Departamento de Matemática

Universidade Federal da Paraíba

João Pessoa, Brazil

Abstract

The notion of mid p -summable sequences was introduced by Karn and Sinha in 2014 and recently explored by Botelho *et al.* in 2017. From this we design a theory of mid summable sequences in the anisotropic setting. As a particular case of our results, we prove that mid p -summable sequences are mid q -summable whenever $p \leq q$, an inclusion result that seems to have been yet not proven in the literature. We will also draw attention to a certain bilinear nature of operators that transform sequences from/into this kind of space.

References

- [1] Botelho, G. and Campos, J. R. *On the transformation of vector-valued sequences by multilinear operators*, Monatsh. Math., **183**, 415–435, 2017.
- [2] Botelho, G., Campos, J. R. and Santos, J. *Operator ideals related to absolutely summing and Cohen strongly summing operators*, Pacific J. of Math., **287**, 1–17, 2017.
- [3] Diestel, J., Jarchow, H. and Tonge, A. *Absolutely Summing Operators*, Cambridge University Press, 1995.
- [4] Karn, A. and Sinha, D. - *An operator summability of sequences in Banach spaces*, Glasg. Math. J., **56**, 427–437, 2014.
- [5] Pietsch, A. *Operator Ideals*, North-Holland, 1980.

Complementation in Fremlin vector lattice symmetric tensor products

Khazhak Navoyan,
Federal University of Uberlândia
Uberlândia, Brazil

Abstract

For a vector lattice E , let $\bar{\otimes}_{n,s}E$ denote the n -fold Fremlin vector lattice symmetric tensor product of E . For $m > n$, we prove that (i) if $\bar{\otimes}_{m,s}E$ is uniformly complete then $\bar{\otimes}_{n,s}E$ is positively isomorphic to a complemented subspace of $\bar{\otimes}_{m,s}E$, and (ii) if there exists $\phi \in E_+^\sim$ such that $\ker(\phi)$ is a projection band in E then $\bar{\otimes}_{n,s}E$ is lattice isomorphic to a projection band of $\bar{\otimes}_{m,s}E$. We also obtain analogous results for the n -fold Fremlin Banach lattice symmetric tensor product $\hat{\otimes}_{n,s,|\pi|}E$ of E where E is a Banach lattice. This is a joint work with professors Dr. Qingying Bu and Dr. Donghai Ji.

References

- [1] F. Blasco, Complementation of symmetric tensor products and polynomials, *Studia Math.* **123** (1997), 165-173.
- [2] Q. Bu, G. Buskes, Polynomials on Banach lattices and positive tensor products, *J. Math. Anal. Appl.* **388** (2012), 845-862.
- [3] S. Dineen, *Complex Analysis on Infinite Dimensional Spaces*, Springer, 1999.
- [4] Kh. Navoyan, Ph.D. dissertation, University of Mississippi, Oxford, Mississippi, USA, 2018.

Biholomorphic Mappings on Banach Spaces

M. Lilian Lourenço

Partially supported by FAPESP 2017/26128-4

e-mail: mllouren@ime.usp.br

Departamento de Matemática- Instituto de Matemática e Estatística

Universidade de São Paulo

Caixa Postal 66281 - CEP : 05315-970 - São Paulo, Brazil

Abstract

H. Cartan theorem stating that a holomorphic self-map of a bounded domain in \mathbb{C}^n with a fixed point at which the derivative is the identity has to be the identity was widened by Cima et al. [2] to separable Hilbert spaces and then to separable dual Banach spaces in [1]. Now, we present an infinite-dimensional version of Cartan theorem concerning the existence of a holomorphic inverse of a given holomorphic self-map of a bounded convex open subset of a dual Banach space. The main assumption is that the derivative operator is power bounded that we, in turn, show to be diagonalizable in some cases, like the separable Hilbert space.

Joint work with H. Carrión and P. Galindo

References

- [1] H. Carrión, P. Galindo and M. L. Lourenço, *Biholomorphic functions in dual Banach spaces*, Complex Anal. Oper. Theory, **(7)**(2013),107-114.
- [2] J. A. Cima, I. Graham, K. T. Kim and S. G. Krantz, *The Caratheodory-Cartan-Kaup-Wu theorem on an infinite dimensional Hilbert space*, Nagoya Math. J. Vol. 185 (2007), 17-30.

Mean Li-Yorke Chaotic Operators on Banach Spaces

Nilson C. Bernardes Jr.

Partially supported by grant #2017/22588-0 (FAPESP), and by CNPq

e-mail: ncbernardesjr@gmail.com

Departamento de Matemática Aplicada

Universidade Federal do Rio de Janeiro

Rio de Janeiro, Brazil

Abstract

We investigate the notion of mean Li-Yorke chaos for operators on Banach spaces. We show that it differs from the notion of distributional chaos of type 2, contrary to what happens in the context of topological dynamics on compact metric spaces. We prove that an operator is mean Li-Yorke chaotic if and only if it has an absolutely mean irregular vector. As a consequence, absolutely Cesàro bounded operators are never mean Li-Yorke chaotic. Dense mean Li-Yorke chaos is shown to be equivalent to the existence of a dense (or residual) set of absolutely mean irregular vectors. As a consequence, every mean Li-Yorke chaotic operator is densely mean Li-Yorke chaotic on some infinite-dimensional closed invariant subspace. A (Dense) Mean Li-Yorke Chaos Criterion and a sufficient condition for the existence of a dense absolutely mean irregular manifold are also obtained. Moreover, several examples and counterexamples are presented. Finally, mean Li-Yorke chaos is also investigated for C_0 -semigroups of operators on Banach spaces.

Joint work with Antonio Bonilla and Alfredo Peris.

References

- [1] N. C. Bernardes Jr., A. Bonilla and A. Peris. *Mean Li-Yorke chaos in Banach spaces*. To appear.

Dynamics of homogeneous polynomials on Banach spaces

Santiago Muro

e-mail: muro@cifasis-conicet.gov.ar

CIFASIS-CONICET

Rosario, Argentina

Rodrigo Cardeccia

e-mail: rcareccia@dm.uba.ar

Departamento de Matemática

Universidad de Buenos Aires

Argentina

Abstract

Let X be a Banach space. A function $F : X \rightarrow X$ is said to be hypercyclic if there exists $x \in X$ whose orbit $Orb_F(x) = \{F^n(x) : n \in \mathbb{N}_0\}$ is dense in X . It is a known fact that there exist hypercyclic linear operators in arbitrary separable infinite dimensional Banach spaces. The dynamical system induced by a (non linear) homogeneous polynomial is quite different. Associated to each homogeneous polynomial there is a ball, centered at zero, with the following property: orbits that meet this limit ball tend to zero. Therefore homogeneous polynomials on Banach spaces cannot be hypercyclic. However, the behavior of the orbits that never enter the limit ball can be non trivial. Indeed, in [1] Bernardes showed the existence of orbits oscillating between infinity and the limit ball. He also proved that there are supercyclic homogeneous polynomials in arbitrary separable infinite dimensional Banach spaces.

In this talk we will exhibit a simple and natural 2-homogeneous polynomial that is at the same time d -hypercyclic (the orbit meets every ball of radius d), weakly hypercyclic (the orbit is dense with respect to the weak topology) and Γ -supercyclic ($\Gamma \cdot Orb_P(x)$ is dense) for each subset $\Gamma \subseteq \mathbb{C}$ unbounded or not bounded away from zero. To prove this, the properties of its Julia set are studied. We will also generalize the construction to arbitrary infinite dimensional Fréchet spaces.

References

- [1] N.C. Bernardes. On orbits of polynomial maps in Banach spaces. *Quaest. Math.*, 21(3-4):318,1998.

Holomorphic functions with distinguished properties on infinite dimensional spaces

Thiago R. Alves

Partially supported by PDJ Ciência sem Fronteiras CNPq Grant 50756/2015-1

e-mail: tralves.math@gmail.com

Departamento de Matemática

Universidade Federal do Amazonas

Manaus, Brazil

G. Botelho

Partially supported by Grant 305958/2014-3 and Fapemig Grant PPM-00450-17

e-mail: botelho@ufu.br

Faculdade de Matemática

Universidade Federal de Uberlândia

Uberlândia, Brazil

Abstract

The study of algebras of holomorphic functions is a classical topic in function theory. When the subject comes to holomorphic functions of infinitely many variables, new phenomena occur, in the sense that functions enjoying properties that are forbidden in the case of finitely many variables might exist (see, e.g, [2]). A central question is the existence, or not, of such functions and, in the positive case, it is usually a very difficult task to construct such functions. A cornerstone in this study was the construction, by Aron, Cole and Gamelin [1], of a bounded holomorphic function on the open unit ball of an infinite dimensional complex Banach space that is continuously but not uniformly continuously extended to the closed unit ball (let us call such functions *Aron-Cole-Gamelin functions*). The main purpose of our talk is to present a method to construct holomorphic functions of infinitely many variables satisfying certain prescribed distinguished properties. For instance, we show how to construct, for the first time to the best of our knowledge, Aron-Cole-Gamelin functions on certain open sets not necessarily the open unit ball. The technique we develop is powerful enough to provide, in certain cases, large algebraic structures formed by such functions (up to the null function, of course).

References

- [1] R. M. Aron, B. J. Cole and T. W. Gamelin, The spectra of algebras of analytic functions on a Banach space, *J. Reine angew. Math.* **415** (1991), 51-93.
- [2] S. Dineen, *Complex Analysis on Infinite Dimensional Spaces*, Springer, 1999.

Positive definite kernels on two-point homogeneous spaces

Victor Simões Barbosa
e-mail: `victor.sb@ufsc.br`
Centro Tecnológico de Joinville
Universidade Federal de Santa Catarina
Joinville-SC, Brasil

Abstract

In this work we study continuous kernels on compact two-point homogeneous spaces which are positive definite and zonal (isotropic). Such kernels were characterized by R. Gangolli some forty years ago and are very useful for solving scattered data interpolation problems on the spaces. In the case the space is the d -dimensional unit sphere, J. Ziegel showed in 2013 that the radial part of a continuous positive definite and zonal kernel is continuously differentiable up to order $\lfloor (d-1)/2 \rfloor$ in the interior of its domain. The main issue here is to obtain a similar result for all the other compact two-point homogeneous spaces.

References

- [1] V.S. Barbosa; V.A. Menegatto. *Differentiable positive definite functions on two-point homogeneous spaces*. J. Math. Anal. Appl. 434 (2016), no. 1, 698-712.
- [2] R. Gangolli. *Positive definite kernels on homogeneous spaces and certain stochastic processes related to Lvy's Brownian motion of several parameters*. Ann. Inst. H. Poincar Sect. B (N.S.) 3 (1967), 121-226.
- [3] J. Ziegel. *Convolution roots and differentiability of isotropic positive definite functions on spheres*. Proc. Amer. Math. Soc. 142 (2014), no. 6, 2063-2077.

Elliptic Partial Differential Equations

Organizer: Edcarlos Domingos da Silva (UFG)

Sharp Regularity for the Inhomogenous Porous Medium Equation

Damião J. Araújo
Partially supported by CNPq
e-mail: araujo@mat.ufpb.br
Departament of Mathematics
Federal University of Paraíba
João Pessoa, Brazil

Abstract

In this talk we shall consider the inhomogeneous porous medium equation

$$\partial u_t - \Delta u^m = f \in L^{q,r} \quad m > 1.$$

Here we show that weak solutions are Hölder continuous, with the following sharp exponent

$$\min \left\{ \frac{\alpha_0^-}{m}, \frac{[(2q - n)r - 2q]}{q[mr - (m - 1)]} \right\},$$

where α_0 denotes the optimal Hölder exponent for solutions of the Homogeneous equation. The method relies on an approximation lemma and geometric iteration with the appropriate intrinsic scaling. This talk is based on *joint* work with J. Miguel Urbano and A.F. Maia - University of Coimbra - Portugal.

Up to the boundary gradient estimates in free boundary problems

Diego Moreira
email: dmoreira@mat.ufc.br
Departament of Mathematics
Universidade Federal do Ceará
Fortaleza, Brazil

Abstract

In this talk, we present some ingredients that lead to gradient estimates up to the boundary in nonlinear free boundary problems. As an application and main motivation, we apply these estimates for singular perturbation problems of flame propagation type.

On nonquadratic fractional coupled elliptic systems in unbounded domains

Edcarlos D. da Silva
IME, UFG
email: edcarlos@ufg.br

Abstract

In this talk we establish existence of positive solutions for a class of nonlocal linearly coupled systems involving Schrödinger equations with fractional Laplacian operator given by

$$\begin{cases} (-\Delta)^{s_1}u + V_1(x)u = f_1(u) + \lambda(x)v, & x \in \mathbb{R}^N, \\ (-\Delta)^{s_2}v + V_2(x)v = f_2(v) + \lambda(x)u, & x \in \mathbb{R}^N, \end{cases}$$

where $(-\Delta)^s$ denotes the fractional Laplacian, $s_1, s_2 \in (0, 1)$ and $N \geq 2$. The coupling function $\lambda : \mathbb{R}^N \rightarrow \mathbb{R}$ is a bounded and continuous function which is related with the potentials by $|\lambda(x)| \leq \delta \sqrt{V_1(x)V_2(x)}$, for some $\delta \in (0, 1)$. We deal with periodic and asymptotically periodic bounded potentials $V_1(x)$ and $V_2(x)$. On the nonlinear terms f_1 and f_2 , we assume “superlinear” at infinity and at the origin. Employing a variational approach we obtain existence of bound and ground states solutions without assuming the well known Ambrosetti-Rabinowitz condition on the nonlinear terms. Furthermore, we give a description of the ground states when the coupling function $\lambda(x)$ goes to zero in the L^∞ norm.

Joint work with João Marcos do Ó (Unb), José Carlos de Albuquerque (UFG).

Hénon Elliptic Equations in \mathbb{R}^2 with Subcritical and Critical Exponential Growth: Mountain–Pass case

Eudes Mendes Barboza
e-mail: eudesmendesbarboza@gmail.com
Departament of Mathematics
Rural Federal University of Pernambuco
Recife, Brasil

João Marcos Bezerra do Ó
e-mail: jmbo.pq@gmail.com
Departament of Mathematics
University of Brasília
Brasília, Brasil

Abstract

We study the Dirichlet problem in the unit ball B_1 of \mathbb{R}^2 for the Hénon-type equation of the form

$$\begin{cases} -\Delta u = \lambda u + |x|^\alpha f(u) & \text{in } B_1, \\ u = 0 & \text{on } \partial B_1, \end{cases}$$

where $f(t)$ is a C^1 -function in the critical growth range motivated by the celebrated Trudinger-Moser inequality. We consider $0 < \lambda < \lambda_1$ and under suitable hypotheses on constant and $f(t)$, by variational methods, we study the solvability of this problem in appropriate Sobolev Spaces.

References

- [1] Adimurthi, *Existence of positive Solutions of the semilinear Dirichlet Problem with critical growth for the n -Laplacian*, Ann. Sc. Norm. Sup. Pisa Cl. Sci. **17** (1990) 393-413.
- [2] A. Ambrosetti, P. H. Rabinowitz, *Dual variational methods in critical point theory and applications*, J. Funct. Anal. **14** (1973) 349-381.
- [3] M. Badiale, E. Serra, *Multiplicity results for the supercritical Hénon equation*, Adv. Nonlinear Stud. **4** (2004) 453-467.
- [4] V. Barutello, S. Secchi, E. Serra, *A note on the radial solutions for the supercritical Hénon equation*, J. Math. Anal. Appl. **341** (2008) 720-728.
- [5] H. Brezis, L. Nirenberg, *Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents*, Comm. Pure Appl. Math. **36** (1983) 437-477.
- [6] H. Brezis, L. Nirenberg, *Remarks on finding critical points*, Comm. Pure and Appl. Math. **54** (1991) 939-963.
- [7] D. Bonheure, E. Serra, M. Tarallo, *Symmetry of extremal functions in Moser-Trudinger inequalities and a Hénon type problem in dimension two*, Adv. Diff. Eq. **13** (2008) 105-138.
- [8] M. Calanchi, E. Terraneo, *Non-radial maximizers for functionals with exponential non-linearity in \mathbb{R}^2* , Adv. Nonlinear Stud. **5** (2005) 337-350.

- [9] P. Carrião, D. de Figueiredo, O. Miyagaki, *Quasilinear elliptic equations of the Hénon-type: existence of non-radial solution*, Commun. Contemp. Math. **11** (2009) 783-798.
- [10] G. Chen, W. Ni and J. Zhou, *Algorithms and visualization for solutions of nonlinear elliptic equations*, Internat. J. Bifur. Chaos Appl. Sci. Engrg. **10** (2000) 1565-1612.
- [11] D. de Figueiredo, P. Lions, R. D. Nussbaum, *A priori estimates and existence of positive solutions of semilinear elliptic equations*, J. Math. Pure Appl. **61** (1982) 41-63.
- [12] D. Figueiredo, O. Miyagaki, B. Ruf, *Elliptic equations \mathbb{R}^2 with nonlinearities in the critical growth range*, Calc. Var. and PDEs. **3** (1995) 139-153.
- [13] D. Figueiredo, J. do Ó, B. Ruf, *On an inequality by N. Trudinger and J. Moser and related elliptic equations*, Comm. Pure Appl. and Math. **55** (2002) 135-152.
- [14] D. de Figueiredo, I. Peral, J. Rossi, *The critical hyperbola for a Hamiltonian elliptic system with weights*, Ann. Mat. Pura Appl. **187** (2008) 531-545.
- [15] M. Hénon, *Numerical experiments on the stability of spherical stellar systems*, Astronomy and Astrophysics **24** (1973) 229-238.
- [16] S. Li, S. Peng, *Asymptotic behavior on the Hénon equation with supercritical exponent*, Sci. China Ser. A. **52** (2009) 2185-2194.
- [17] P. Lions, *The concentration-compactness principle in the calculus of variations The limit case I*, Revista Iberoamericana **1** (1985) 145-201.
- [18] W. Long, J. Yang, *Existence for critical Hénon-type equations*, Diff. and Int. Eq. **25** (2012) 567-578.
- [19] W. Long, J. Yang, *Existence of solutions for critical growth Hénon equations*, J. Jiangxi Normal Uni. **5** (2010) 463-466.(in Chinese)
- [20] J. Moser, *A sharp form of an inequality by N. Trudinger*, Ind. Univ. Math. J. **20** (1971) 1077-1092.
- [21] W. Ni, *A nonlinear Dirichlet problem on the unit ball and its applications*, Ind. Univ. Math. J. **31** (1982) 801-807.
- [22] S. Peng, *Multiple boundary concentrating solutions to Dirichlet problem of Hénon equation*, Acta Math. Appl. Sin. Engl. Ser. **22** (2006) 137-162.
- [23] P. H. Rabinowitz, *Minimax Methods in Critical Point Theory with Applications to Differential Equations*, CBMS Reg. Conf. Ser. Math. 65, Amer. Math. Soc., Providence, RI, 1986.
- [24] P.H. Rabinowitz, *On a class of nonlinear Schrödinger equations*, Z. Angew Math. Phys. **43** (1992) 270-291.
- [25] B. Ribeiro, *Critical elliptic problems in \mathbb{R}^2 involving resonance in high-order eigenvalues*, Commun. Contemp. Math. **17** (2015) 22 pp.
- [26] S. Secchi, E. Serra, *Symmetry breaking results for problems with exponential growth in the unit disk*, Commun. Contemp. Math. **8** (2006) 823-839.
- [27] E. Serra, *Nonradial positive solutions for the Hénon equation with critical growth*, Calc. Var. and PDEs. **23** (2005) 301-326.

- [28] E. A. de B. e Silva, *Linking theorems and applications to semilinear elliptic problems at resonance*, N. Anal., Theory, Meth. Appl. **16** (1991) 455-477.
- [29] D. Smets, J. Su, M. Willem, *Non-radial ground states for the Hénon equation*, Commun. Contemp. Math. **4** (2002) 467-480.
- [30] C. Tarsi, *On the existence and radial symmetry of maximizers for functionals with critical exponential growth in \mathbb{R}^2* , Diff. Int. Eq. **21** (2008) 477-495.
- [31] N. Trudinger, *On imbedding into Orlicz spaces and some applications*, J. Math. Mech. **17** (1967) 473-483.

Infinitely many solutions for a class of degenerated or singular elliptic p -Kirchhoff equations

Gustavo Ferron Madeira
e-mail: gfmadeira@dm.ufscar.br
Departament of Mathematics
Federal University of São Carlos
São Carlos–SP, Brasil

Abstract

We are concerned in this lecture with existence of infinitely many solutions for a class of elliptic equations where Kirchhoff term may be degenerated, discontinuous or singular at origin. The source term in the equation may be nonlocal as well. In the final part of the lecture extension of the results to more general nonhomogeneous elliptic Kirchhoff equations will be also discussed.

Least energy nodal solution for the Schrödinger-Poisson system under an asymptotically cubic nonlinearity

Edwin G. Murcia
e-mail: murciae@javeriana.edu.co
Departamento de Matemáticas
Pontificia Universidad Javeriana
Bogotá, Colombia

Gaetano Siciliano
Partially supported by Capes, CNPq e Fapesp
e-mail: sicilian@ime.usp.br
Departamento de Matemática
Universidade de São Paulo
São Paulo, Brazil

Abstract

In this talk we present a recent result concerning the existence of a least energy sign changing solution for a nonlinear Schrödinger-Poisson system in the whole space. The nonlinearity behaves asymptotically cubic at infinity then it is in strict competition with the nonlocal term appearing in the energy functional. To prove the existence of such a solution we work on the the "nodal Nehari set" on which we minimize the functional. One of the main difficulties is to show that this set is nonempty under the assumption made on the nonlinearity.

References

- [1] E.G. Murcia, G. Siciliano, *Least energy radial sign-changing solution for the Schrödinger - Poisson system in \mathbb{R}^3 under an asymptotically cubic nonlinearity*, *arXiv:1805.00259*

Extremal for a k -Hessian inequality of Trudinger-Moser type

José Francisco de Oliveira
 e-mail: jfoliveira@ufpi.edu.br
 Federal University of Piauí
 Teresina, Brazil

João Marcos do Ó
 Brasília University
 Brasília, Brazil

Bernhard Ruf
 Università di Milano
 Milan, Itália

Abstract

We consider k -Hessian operators $S_k[u]$ in bounded domains Ω in \mathbb{R}^N such that $\partial\Omega$ is $(k-1)$ -convex. For so-called k -admissible functions $u \in \Phi_0^k$ one has Sobolev type inequalities of the form

$$\|u\|_{L^p(\Omega)} \leq C \|u\|_{\Phi_0^k}$$

where $\|u\|_{\Phi_0^k}^{k+1} = \int_{\Omega} (-u) S_k[u] dx$, and $1 \leq p \leq k^* = \frac{N(k+1)}{N-2k}$. The case $N = 2k$ is a borderline case of Trudinger-Moser type, and it was recently shown by Tian-Wang that a corresponding inequality of exponential type holds

$$\sup_{\|u\|_{\Phi_0^k} \leq 1} \int_{\Omega} \left(e^{\alpha|u|^{\frac{N+2}{N}}} - \sum_{j=0}^{k-1} \frac{\alpha^j |u|^{j \frac{N+2}{N}}}{j!} \right) dx \leq C$$

for $\alpha \leq \alpha_N = N \left[\frac{\omega_{N-1}}{k} \binom{N-1}{k-1} \right]^{2/N}$. In this article we prove an analogue to the famous result of Carleson–Chang, namely that for $\Omega = B_R(0)$ the above supremum is attained also in the limiting case $\alpha = \alpha_N$.

References

- [1] L. Carleson, S. Y. A. Chang, *On the existence of an extremal function for an inequality of J. Moser*, Bull. Sci. Math. **110** (1986), 113–127.
- [2] J. Moser, *A sharp form of an inequality by N. Trudinger*, Indiana Univ. Math. J. **20** (1970/71), 1077–1092.
- [3] G. -T, Tian, X. -J. Wang, *Moser-Trudinger type inequalities for the Hessian equation*, J. Funct. Anal. **259** (2010) 1974–2002

On the well-posedness of Goursat problems in Gevrey classes

Jorge Marques

Partially supported by CeBER and FEUC

e-mail: jmarques@fe.uc.pt

School of Economics

University of Coimbra

Coimbra, Portugal

Abstract

Some authors, Nishitani [5], Hasegawa [3], Carvalho e Silva [4], have investigated the C^∞ well-posedness of Goursat problems for linear PDE's with constant coefficients. I am interested in trying to find necessary and sufficient conditions for the generalized Goursat problem to be well-posed in the Gevrey classes Γ^s with $s > 1$.

References

- [1] Gevrey, M.: Sur la nature analytique des solutions des équations aux dérivées partielles. *Ann. Ecole Norm. Sup. Paris*, **35**, 129–190, 1918.
- [2] Hadamard, J.: *Le problème de Cauchy et les équations aux dérivées partielles linéaires hyperboliques*. Hermann, Paris, 1932.
- [3] Hasegawa, Y.: On the C^∞ Goursat problem for equations with constant coefficients. *J. Math. Kyoto Univ.*, **19**, 125–151, 1979.
- [4] Marques, J., Carvalho e Silva, J.: Gevrey well posedness of Goursat-Darboux problems and asymptotic solutions. In Pinelas, S., Caraballo, T., Kloeden, P., Graef, J.R. (Eds.). *Differential and Difference Equations and Applications*. Springer International Publishing, 2018.
- [5] Nishitani, T.: On the C^∞ well-posedness for the Goursat problem with constant coefficients, *J. Math. Kyoto Univ.*, **20**, 179–190, 1980.

The fibering method approach for a non-linear Schrödinger equation coupled with the electromagnetic field

Kaye Silva

e-mail: kayeoliveira@hotmail.com, kaye.Oliveira@ufg.br

Instituto de Matemática e Estatística

Universidade Federal de Goiás

Goiânia, Brazil

Gaetano Siciliano

e-mail: sicilian@ime.usp.br

Departamento de Matemática

Universidade de São Paulo

São Paulo, Brazil

Abstract

We study, with respect to the parameter $q \neq 0$, the following Schrödinger-Bopp-Podolsky system in \mathbb{R}^3

$$\begin{cases} -\Delta u + \omega u + q^2 \phi u = |u|^{p-2}u, \\ -\Delta \phi + a^2 \Delta^2 \phi = 4\pi u^2, \end{cases}$$

where $p \in (2, 3]$, $\omega > 0$, $a \geq 0$ are fixed. We prove, by means of the fibering approach, that the system has no solutions *at all* for large values of q 's, and has two *radial* solutions for small q 's (see [1]). We give also qualitative properties about the energy level of the solutions and a variational characterization of these extremal values of q . Our results recover and improve some results in [2, 3].

References

- [1] Y. Ilyasov. *On extreme values of Nehari manifold method via nonlinear Rayleigh's quotient*. Topological Meth. Nonlinear Anal. **49** (2017) 683–714.
- [2] d'Avenia, Pietro and Siciliano, Gaetano. *Nonlinear Schrödinger equation in the Bopp-Podolsky electrostatics: solutions in the electrostatic case*. arXiv:1802.03380 (2018).
- [3] Ruiz, David. *The Schrödinger-Poisson equation under the effect of a nonlinear local term*. J. Diff. Equations **237** (2006), 655–674.

Ground states for quasilinear Schrödinger elliptic systems

Maxwell L. Silva

A joint work with **Edcarlos D. Silva** and **J.C. de Albuquerque**

e-mail: maxwelllizete@hotmail.com

Institute of Mathematics and Statistics

Federal University of Goiás

Goinia, Brazil

Abstract

In this work we are concerned with the existence and nonexistence of ground state solutions for the following class of quasilinear Schrödinger coupled systems

$$\begin{cases} -\Delta u + a(x)u - \Delta(u^2)u = g(u) + \theta\lambda(x)uv^2, & x \in \mathbb{R}^N, \\ -\Delta v + b(x)v - \Delta(v^2)v = h(v) + \theta\lambda(x)vu^2, & x \in \mathbb{R}^N, \end{cases}$$

where $N \geq 3$, $\theta \geq 0$, $a, b, \lambda : \mathbb{R}^N \rightarrow \mathbb{R}$ are periodic or asymptotically periodic functions. The nonlinear terms g, h are superlinear at infinity and at the origin. By using a change of variable, we turn the quasilinear system into a nonlinear system where we can establish a variational approach with a fine analysis on the Nehari method. For the nonexistence result we compare the potentials $a(x), b(x)$ with periodic potentials proving nonexistence of ground state solutions.

References

- [1] A. Ambrosetti, E. Colorado, *Standing waves of some coupled nonlinear Schrödinger equations*, J. Lond. Math. Soc. (2) 75 (2007), no. 1, 67-82.
- [2] Y. Guo, Z. Tang, *Ground state solutions for quasilinear Schrödinger systems*, J. Math. Anal. Appl. 389 (2012), no. 1, 322-339.
- [3] L. A. Maia, E. Montefusco, B. Pellacci, *Weakly coupled nonlinear Schrödinger systems: the saturation effect*, Calc. Var. Partial Differential Equations 46 (2013), no. 1-2, 325-351.
- [4] L. A. Maia, E. Montefusco, B. Pellacci, *Positive solutions for a weakly coupled nonlinear Schrödinger system*, J. Differential Equations 229 (2006), no. 2, 743-767.
- [5] E. A. B. Silva, G. F. Vieira, *Quasilinear asymptotically periodic Schrödinger equations with subcritical growth*, Nonlinear Analysis 72 (2010) 2935-2949.
- [6] E. D. Silva, M. L. Silva, J. C. de Albuquerque, *Positive ground states solutions for a class of quasilinear coupled superlinear elliptic systems*, to appear.
- [7] Y. Shen, Y. Wang, *Soliton solutions for generalized quasilinear Schrödinger equations*, Nonlinear Analysis 80 (2013) 194-201.
- [8] M. A. S. Souto, S. H. M. Soares *Ground state solutions for quasilinear stationary Schrödinger equations with critical growth*, Commun. Pure Appl. Anal. 12 no 1 (2013) 99-116.

On the Choquard equation with Hardy-Littlewood-Sobolev upper critical exponent

Minbo Yang

Partially supported by NSFC(11571317)

e-mail: mbyang@zjnu.edu.cn

Department of Mathematics

Zhejiang Normal University

Jinhua, P.R. China

Abstract

In this talk I am going to introduce some existence results for the Choquard equation

$$-\varepsilon^2 \Delta u + V(x)u = \varepsilon^{\mu-3} \left(\int_{R^N} \frac{Q(y)G(u(y))}{|x-y|^\mu} dy \right) Q(x)g(u) \quad \text{in } R^N,$$

where $0 < \mu < N$, ε is a positive parameter, V, Q are two continuous real function on R^N and G is the primitive of g which is of critical growth due to the Hardy-Littlewood-Sobolev inequality. under different assumptions on the nonlinearities and potentials, we proved the existence and multiplicity of solutions, including the Brezis-Nirenberg type results, for this nonlocal equation by variational methods.

References

- [1] C. O. Alves, F. Gao, M. Squassina and M. Yang. Singularly perturbed critical Choquard equations, *J. Differential Equations*, **263** (2017), 3943–3988.
- [2] H. Brézis and L. Nirenberg. Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents, *Comm. Pure Appl. Math.*, 36 (1983), 437–477.
- [3] F. Gao and M. Yang. On the Brezis-Nirenberg type critical problem for nonlinear Choquard equation, *Sci China Math.*, doi: 10.1007/s11425-016-9067-5
- [4] F. Gao and M. Yang. A strongly indefinite Choquard equation with critical exponent due to Hardy-Littlewood-Sobolev inequality, *Commun. Contemp. Math.*, <https://doi.org/10.1142/S0219199717500377>.

Super critical problems with concave and convex nonlinearities in \mathbb{R}^N

J. M. do Ó

e-mail: jmbo@pq.cnpq.br
Department of Mathematics
Brasília University
Brasília, DF, Brazil

P. K. Mishra

Partially supported by PNPd-CAPES
e-mail: pawanmishra@mat.ufpb.br
Department of Mathematics
Federal University of Paraíba
João Pessoa-PB, Brazil

A. Moameni

e-mail: momeni@math.carleton.ca
School of Mathematics and Statistics
Carleton University
Ottawa, Ontario, Canada

Abstract

In this work, adopting a methodology based on variational principle on convex sets, we establish an existence and multiplicity result for a class of semilinear elliptic problems defined on whole \mathbb{R}^N with nonlinearities involving sublinear and superlinear terms.

References

- [1] Antonio Ambrosetti, Haïm Brezis, and Giovanna Cerami. *Combined effects of concave and convex nonlinearities in some elliptic problems*. J. Funct. Anal., **122**(2), 519–543, 1994.
- [2] J. Chabrowski and João Marcos Bezzera do Ó. *On semilinear elliptic equations involving concave and convex nonlinearities*. Math. Nachr., **233/234**, 55–76, 2002.
- [3] Abbas Moameni. *A variational principle for problems with a hint of convexity*. C. R. Math. Acad. Sci. Paris, **355**(12), 1236–1241, 2017.
- [4] Andrzej Szulkin. *Minimax principles for lower semicontinuous functions and applications to nonlinear boundary value problems*. Ann. Inst. H. Poincaré Anal. Non Linéaire, **3**(2), 77–109, 1986.

Parabolic Differential Equations

Organizer: Anderson Luis Albuquerque de Araújo (UFV)

The number π and the nonexistence of solution for Chern-Simons equation

Adilson Eduardo Presoto

Partially supported by FAPESP and CAPES

e-mail: `presoto@dm.ufscar.br`

Departament of Mathematics

Federal University of São Carlos

São Carlos, Brazil

Abstract

The Brezis and Benilan's seminal work [1] on Thomas-Fermi problem which leads with elliptic problems involving L^1 or measure data has guided us to a new class of fascinating problems. Here we are interesting in understand the phenomenon of breaking of existence solutions of

$$\begin{cases} -\Delta u + e^u(e^u + 1) = \mu, & \text{in } \Omega, \\ u = 0, & \text{on } \partial\Omega. \end{cases}$$

when the measure evaluated in singletons surpass $4\pi\mathcal{H}^{n-2}$. We approximate the datum by a suitable convergent sequence in the measure sense in order to obtain a convergent sequence of solutions. The limit will be determined in terms only of datum. We also discuss the unlike cases when the data has signal and the counterpart of system.

References

- [1] PH. BÉNILAN; H. BRÉZIS, Nonlinear problems related to the Thomas-Fermi equation. *J. Evol. Equ.* **3** (2003), no. 4, 673–770.
- [2] H. BREZIS; M. MARCUS; A. C. PONCE, Nonlinear elliptic equations with measure revisited. In J. Bourgain, C. E. Kenig, and S. Klainerman (eds.), *Mathematical aspects of nonlinear dispersive equations*. Ann. of Math Studies 163, Princeton University Press, Princeton, NJ, 2007, 55–109.
- [3] A. C. PONCE; A. E. PRESOTO, Limit solutions of the Chern-Simons equation. *Nonlinear Anal.* **84** (2013), 91–102.

A class of parabolic equations driven by the mean curvature flow

Anderson L. A. de Araujo

Partially supported by FAPESP 2013/22328 – 8

e-mail: anderson.araujo@ufv.br

Departament of Mathematics

Federal University of Viçosa

Viçosa - MG, Brazil

Marcelo Montenegro

Partially supported by by CNPq

e-mail: msm@ime.unicamp.br

Department of Mathematics

State University of Campinas

Campinas - SP, Brazil

Abstract

We study a class of parabolic equations which can be viewed as a generalized mean curvature flow acting on cylindrically symmetric surfaces with Dirichlet condition on the boundary. We prove existence of a unique solution by means of an approximation scheme. We also develop the theory of asymptotic stability for solutions of general parabolic problems.

References

- [1] A.L.A. de Araujo, M. S. Montenegro. *A class of parabolic equations driven by the mean curvature flow*. To appear in Proceedings of the Edinburgh Mathematical Society, 2018.

Global solutions for a fractional reaction–diffusion equation

Arlúcio Viana
e-mail: arlucioviana@gmail.com
Departament of Mathematics
University of Sergipe
Itabaiana, Brazil

Abstract

We shall discuss the existence and nonexistence of global positive solutions for the semilinear fractional diffusion equation

$$u_t(t, x) = \int_0^t dg_\alpha(s) \Delta u(t - s, x) + |u(t, x)|^{\rho-1} u(t, x), \text{ in } (0, \infty) \times \mathbb{R}^N; \quad (3)$$

$$u(x, 0) = u_0(x), \text{ in } \mathbb{R}^N, \quad (4)$$

where $\rho > 1$. Indeed, a combination of [1, Th. 1] and [2, Th. 5] provides the following result:

- (i) If $1 < \rho < 1 + \frac{2}{\alpha N}$, there exists no nonnegative global solution of (3)-(4).
- (ii) If $\rho \geq 1 + \frac{2}{\alpha N}$, there exists a global nonnegative solution of (3)-(4).

Eventually, we highlight some asymptotic properties of the existing global solution.

References

- [1] B. de Andrade and A. Viana, On a fractional reaction-diffusion equation, *Z. Angew. Math. Phys.* **68** (2017), no. 3, Art. 59, 11 pp.
- [2] A. Viana, A local theory for a fractional reaction-diffusion equation, *Comm. Cont. Math.*, doi: 10.1142/S0219199718500335.

On the Dirichlet-to-Neumann Semigroup: Recent Developments and Results

Jamil Abreu

E-mail: jamil.abreu@ufes.br

Department of Applied Mathematics

Federal University of Espírito Santo

São Mateus - ES, Brazil

Abstract

Let $\Omega \subset \mathbb{R}^N$ be an open set with Lipschitz boundary. The Dirichlet-to-Neumann operator, in its simplest version, is the operator D_0 acting on $L^2(\partial\Omega)$ with the property that $\varphi \in \mathcal{D}(D_0)$ and $D_0\varphi = h$ if, and only if there exists $u \in H^1(\Omega)$ with $\Delta u = 0$ on Ω , $u|_{\partial\Omega} = \varphi$ on $\partial\Omega$, and $\partial_\nu u = h$ in a weak sense. It turns out that D_0 is the associated operator to a j -elliptic form (following Arendt and ter Elst [2]), namely, the classical Dirichlet form

$$a(u, v) = \int_{\Omega} \nabla u \cdot \overline{\nabla v} dx,$$

where $j : H^1(\Omega) \rightarrow L^2(\partial\Omega)$ is the trace operator from $H^1(\Omega)$ to $L^2(\partial\Omega)$.

In this talk we comment on several results concerning some other versions of this operator, including the case of rough domains and more general dynamics such as $\mathcal{L}u = \lambda u$, where \mathcal{L} is a general second order differential operator satisfying certain hypotheses.

References

- [1] J. Abreu and É. Capelato. Dirichlet-to-Neumann semigroup with respect to a general second order eigenvalue problem. *Semigroup Forum* (to appear).
- [2] W. Arendt and A.F.M. ter Elst. Sectorial forms and degenerate differential operators. *J. Operator Theory*, (2012) **67** 33-72.

Fractional Regularity for Degenerate Equations

Luís H. de Miranda

Partially supported by FAPDF - Brazil

e-mail: demiranda@unb.br

Department of Mathematics

Universidade de Brasília

Brasília, Brazil

Abstract

In this talk we are going to address the fractional regularity of solutions for Quasilinear degenerate equations of both parabolic and elliptic type. Our goal will be to link the interaction between the nonlinear character of p -Laplacian-like operators with certain spaces of fractional order of differentiability and its effects on the associated solutions. In turn, special attention will be delivered for the case of the parabolic p -Laplacian, cf. [1], and also to the (p, q) -Laplacian, cf. [2], where we are going to review or present some new and old results regarding the regularity of solutions to this sort of equations, as well as the related a priori estimates.

References

- [1] L.H. de Miranda and G. Planas. Parabolic p -Laplacian revisited: global regularity and fractional smoothness. *Commun. Contemp. Math.*, *to appear* (2018)
- [2] L.H. de Miranda and A. Presoto. On the fractional regularity for degenerate equations with (p, q) -growth. Preprint (2018)

Attractors for a vectorial von Karman system

Rodrigo Nunes Monteiro
Partially supported by PNPd/CAPES
e-mail: rodrigonunesmonteiro@gmail.com
Department of Mathematics
Federal University of Viçosa - UFV
Viçosa, Brazil

Abstract

This lecture is concerned with long-time dynamics of a full von Karman (vectorial) system subject to nonlinear thermal coupling and *free* boundary conditions. In contrast with *scalar* von Karman system, vectorial full von Karman system accounts for both vertical and in plane displacements. The corresponding PDE is of critical interest in flow structure interactions where nonlinear plate/shell dynamics interacts with perturbed flows [vicid or invicid]. It is shown that the system is quasi-stable and admits a global attractor which is also *smooth and of finite fractal dimension*. The above result is shown to hold for plates without regularizing effects of rotational inertia and without any mechanical dissipation imposed on vertical displacements.

References

- [1] I. Chueshov and I. Lasiecka, Long-Time Behavior of Second order Evolutions with Nonlinear Damping, Mem. Amer. Math. Soc. 195 (2008), no. 912.
- [2] I. Chueshov and I. Lasiecka, Von Karman Evolution Equations. Well-posedness and Long Time Dynamics, Springer Monographs in Mathematics. Springer, New York, 2010.
- [3] I. Lasiecka and R. Triggiani, *Analyticity of thermo-elastic semigroups with free boundary conditions*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 27 (1998), no. 3-4, 457-482.

Poster Session

Organizer: Willian Cintra (UnB)

The Riemann hypothesis as a problem of functional analysis

Alfredo Sotelo Pejerrey

Partially supported by Pontificia Universidad Católica del Perú

e-mail: sotelo.alfredo@pucp.pe

Departament of Mathematics

Pontificia Universidad Católica del Perú

Lima, Perú

Abstract

J. Alcántara [1] has proved that the Riemann Hypothesis (RH) holds true if and only if a certain compact operator A_ρ on $L^2(0,1)$ is injective. In [2] and [3] has been establish several relations between the operators A_ρ and $A_\rho(\alpha)$. One of then helps us to establish, using Theorem 1 in [4], that RH holds true if and only if $Ran(V_\alpha) \subset \overline{Ran(A_\rho^*(\alpha))}$.

In [1] has been proven that A_ρ is non-nuclear Hilbert-Schmidt. Using the fact that if (λ_n) is the sequence of non-zero eigenvalues of A_ρ where the ordering is such that $|\lambda_n| \geq |\lambda_{n+1}|$ for all $n \in \mathbb{N}$ and each one of them being repeated according to its algebraic multiplicity, then the first eigenvalue λ_1 is positive and has algebraic multiplicity one, $|\lambda_n| \leq \frac{e}{n}$ for all $n \in \mathbb{N}$, we stablish, using the Ringrose decomposition [5] and the spectral theorem for compact normal operators, that the sum of A_ρ and a quasi-nilpotent operator belongs to Lorentz ideal of compact operators $M_{1,\infty}$.

References

- [1] J. Alcántara-Bode. *An integral equation formulation of the Riemann Hypothesis*. J. Integral Equation and Operator Theory, **17**, (1993), 151-168.
- [2] J. Alcántara-Bode. *A completeness problem related to the Riemann Hypothesis*. J. Integral Equation and Operator Theory, **53 (3)**, (2005), 301-309.
- [3] J. Alcántara-Bode. *An Algorithm for the Evaluation of certain Fredholm Determinants*. J. Integral Equation and Operator Theory, **39**, (2001), 153-158.
- [4] R. G. Douglas. *On majorization, factorization, and range inclusion of operators on Hilbert spaces*. Proc. Amer. Math. Soc., **17**, (1966), 413-415.
- [5] J. Ringrose. *Super-diagonal forms for compact linear operators*. Proc. Lond. Math. Soc., **12 (3)**, (1962), 367-384.

Relation between approximation tools in homogeneous compact spaces of rank 1

Angelina Carrijo de Oliveira Ganancin Faria

Partially supported by CAPES

e-mail: angelina.carrijo@usp.br

Instituto de Ciências Matemáticas e de Computação

Universidade de São Paulo

São Carlos, Brasil

Abstract

In constructive approximation theory there are a lot of tools measuring, some how, the smoothness of functions. In this work we are interested on finding relations between particular of these tools, namely, moduli of smoothness, K-functionals and the rate of approximation of average operators on homogeneous compact spaces of rank 1.

It is proved a generalization of a recent result obtained on the spherical setting. Specifically, we show that the rate of approximation of the shifting operator is equivalent (in the asymptotic meaning) to the classical K-functional defined on homogeneous compact spaces of rank 1, and given by Laplace-Beltrami operator.

Relations once established permit us to apply a very new technique in order to get sharp estimates for the decay rate of eigenvalue sequences of certain integral operators on homogeneous compact spaces of rank 1.

References

- [1] A. O. Carrijo; T. Jordo. *Approximation tools on compact two-point homogeneous spaces: characterizations and applications*. 2018. Submitted for publication. <https://arxiv.org/abs/1708.02576>.
- [2] F. Dai; Z. Ditzian. *Combinations of multivariate averages*. J. Approx. Theory, v. 131, n. 2, p. 268-283, 2004.
- [3] S. S. Platonov. *Some problems in the theory of the approximation of functions on compact homogeneous manifolds*. Mat. Sb., v. 200, n. 6, p. 67-108, 2009.

Standing waves for a system of quasilinear schrödinger equations in \mathbb{R}^N

João Marcos do Ó

Partially supported by CNPq, PRONEX-MCT/Brazil and the Millennium Institute for the Global Advancement of Brazilian Mathematics

e-mail: jmbo@pq.cnpq.br
Departament of Mathematics
Federal University of Paraíba
João Pessoa, PB, Brazil

Olimpio Miyagaki

Partially supported by CNPq
e-mail: ohmiyagaki@gmail.com
Departament of Mathematics
Federal University of Juiz de Fora, MG, Brazil

Cláudia Santana

Partially supported by UESC
e-mail: santana@uesc.br
Department of Exact and Technological Sciences
State University of Santa Cruz
Ilhéus, BA, Brazil

Abstract

In this paper we study the existence of bound state solutions for stationary quasilinear Schrödinger systems of the form

$$\begin{cases} -\Delta u + V(x)u - [\Delta(u^2)]u = K(x)H_u(u, v) & \text{in } \mathbb{R}^N, \\ -\Delta v + V(x)v - [\Delta(v^2)]v = K(x)H_v(u, v) & \text{in } \mathbb{R}^N, \end{cases} \quad (S)$$

where $N \geq 3$, V and K are bounded continuous nonnegative functions, and, the primitive of nonlinearity, $H(u, v)$ is a p -homogeneous function of class C^1 with $4 < p < 22^*$. We give a special attention to the case when V may eventually vanishes. Our arguments are based on truncation methods, Moser iteration scheme and some variational arguments like as Mountain-Pass Theorem and changing of variable.

References

- [1] ALVES, C. O. - Local Mountain Pass for a class of elliptic system. *J. Math. Anal. Appl.*, **335**, 135-150, 2007.
- [2] ALVES, C. O. AND SOUTO, M. A. - Existence of solutions f or a class of elliptic equations in \mathbb{R}^N with vanishing potentials. *J. Differential Equations*, **252**, 5555-5568, 2012.
- [3] AMBROSETTI, A. AND WANG, Z.-Q. - Nonlinear Schrödinger equations with vanishing and decaying potentials. *Differential Integral Equations*, **18**, 1321-1332, 2005.
- [4] BRÈZIS, H. AND KATO, T. - Remarks on the Scrödinger operator with regular complex potentials. *J. Math. Pures Appl.*, **58**, 137-151, 1979.

- [5] COLIN, M. AND JEANJEAN, L. - Solutions for a quasilinear Schrödinger equation: a dual approach. *Nonlinear Anal.*, **56**, 212-226, 2004.
- [6] DEL PINO, M. AND FELMER, P. - Local mountain pass for semilinear elliptic problems in unbounded domains. *Calc. Var. Partial Differential Equations*, **4**, 121-137, 1996.
- [7] DO Ó, J. M. AND SEVERO, U. - Quasilinear Schrödinger equations involving concave and convex nonlinearities. *Commun. Pure Appl. Anal.*, **8**, 621-644, 2009.

Standing waves for weakly coupled nonlinear Schrodinger systems with critical growth in \mathbb{R}^N

Claudiney Goulart

Partially supported by CNPq (Grant 620108/2008-8)

e-mail: claudiney@ufg.br

Departament of Mathematics

University of Jataí, Jataí, Brazil

Elves Alves de Barros e Silva

Partially supported by CNPq (Grant 620108/2008-8)

e-mail: elves@unb.br

Department of Mathematics

University of Brasília, Brasília, Brazil

Abstract

In this work we apply variational methods to study the existence of positive solutions for the following nonlinear coupled Schrodinger system with critical growth in \mathbb{R}^N

$$\begin{cases} -\Delta u + \lambda_1 u &= |u|^{p-2}u + \frac{2\beta\alpha}{\alpha+\mu}|u|^{\alpha-2}u|v|^\mu, & \text{in } \mathbb{R}^N, \\ -\Delta v + \lambda_2 v &= |v|^{q-2}v + \frac{2\beta\mu}{\alpha+\mu}|v|^{\mu-2}v|u|^\alpha, & \text{in } \mathbb{R}^N, \end{cases} \quad (5)$$

with $N \geq 2$, $\beta > 0$, $\alpha, \mu > 1$, $\alpha + \mu = 2^*$, $2 < p, q < 2^*$ and $\lambda_1, \lambda_2 > 0$, where we define $2^* = \infty$ if $N = 2$ and $2^* = 2N/(N - 2)$ if $N \geq 3$. Existence of positive least energy solution will also be established. For this, we consider the associated functional restricted to Nehari manifold and we apply local and global minimization arguments combined with minimax methods. We also establish appropriate estimates for the level determined by Mountain Pass Theorem and use arguments developed by Brezis and Nirenberg for the study of semilinear problems with critical growth.

References

- [1] A. AMBROSETTI and E. COLORADO, Bound and ground states of coupled nonlinear Schrodinger equations, *C. R. Math. Acad. Sci. Paris* **342** (2006), 453-458.
- [2] H. BRÉZIS, L. NIRENBERG. Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents. *Comm. Pure Appl. Math.*, **36(4)** (1983) , 437-477.
- [3] L. A. MAIA, E. MONTEFUSCO and B. PELLACCI, Positive solutions for a weakly coupled nonlinear Schrodinger system, *J. Differ. Equ.* **229** (2006), 743-767.

Mountain pass algorithm via Pohozaev manifold

Daniel Raom Santiago
Department of Mechanical Engineering
University of Brasilia
Brasilia, Brazil

L. A. Maia
Department of Mathematics
University of Brasilia
Brasilia, Brazil

R. Ruviano
Department of Mathematics
University of Brasilia
Brasilia, Brazil

Y. D. Sobral
Department of Mathematics
University of Brasilia
Brasilia, Brazil

Abstract

A new numerical algorithm for solving asymptotically linear problems is presented. The ground state solution of the problem, which in general is obtained as a min-max of the associated functional, is obtained as a minimum of the functional constrained to the Pohozaev manifold instead. Examples are given of the use of this method for finding numerical solutions depending on various parameters.

References

- [1] L. Jeanjean, K. Tanaka. *A remark on least energy solutions in \mathbb{R}^N* . Proc. Amer. Math. Soc., 131, no. 8 (2002), 2399-2408
- [2] G. Chen, J. Zhou and Wei-Ming Ni. *Algorithms and visualizations for solutions of nonlinear elliptic equations*. International Journal of Bifurcation and Chaos, Vol. 10, 7 (2000), 1565-1612

On retarded potentials and applications

Felipe B. Federson

Email: felipe.federson@usp.br

Instituto de Física de São Carlos

Universidade de São Paulo, São Carlos, Brazil

Everaldo M. Bonotto

Supported by FAPESP and CNPq

Email: ebonotto@icmc.usp.br

Instituto de Ciências Matemáticas e de Computação

Universidade de São Paulo, São Carlos, Brazil

Abstract

When we take into account time-varying electric current or charge distributions in the past, it is reasonable to consider retarded potentials for the generated electromagnetic fields. The fields propagate at the speed of light, c , so the delay of the fields connecting cause and effect at earlier and later times is an important factor. We extend the notion of retarded potentials to paths so that we can employ the Henstock integral in order to obtain a Feynman-Kač-type formula to the wavefunction which is a solution of the Schrödinger equation with retarded potentials in quantum mechanics.

References

- [1] E.M. Bonotto; M. Federson; P. Muldowney, A Feynman-Kač solution to a random impulsive equation of Schrödinger type, *Real Anal. Exchange* **36**(1), (2010/11), 107-148.
- [2] D. J. Griffiths, *Introduction to Electrodynamics*, Prentiss-Hall, 1989.
- [3] P. Muldowney, *A Modern Theory of Random Variation - The Henstock Integral in Probability and Stochastic Calculus, with application to Option Pricing and Feynman Path Integrals*, John Wiley & Sons, Inc. 2012.

Topological optimization in the kirchhoff plate bending model using topological derivative

Fernando S. Carvalho

e-mail: fscarvalho@uft.edu.br

Department of Mechanical Engineering - Faculty of Technology

University of Brasilia

Brasilia, Brazil

Abstract

The topological derivative is a scalar field that measures the sensitivity of a functional data to an infinitesimal perturbation in a $\Omega \subset \mathbb{R}^n$ domain. The topological derivative has been successfully used for topological optimization in a large class of engineering and physical problems. In this work, we use the topological derivative in the Kirchhoff plate bending model, which is modeled by a fourth-order differential operator, in order to maximize the first eigenvalue, which represents the first natural frequency of vibration.

References

- [1] Amstutz, S.; Novotny, A.A.. *Topological asymptotic analysis of the Kirchhoff plate bending problem*
- [2] Novotny, A. A.; Sokolowisky, J.; *Análise de Sensibilidade Topológica - Teoria e Aplicações*, SBMAC, Coleo: Notas em Matemática Aplicada, So Carlos - SP, 2016.
- [3] Feijo, R. A.; *The Topological Derivative for the Poisson's Problem*. Mathematical Models and Methods in Applied Sciences, v. 13, n. 12, p. 1825-1844, 2003.

Solving D’Alambert problem using Perplex Numbers

Ian Rodrigues Massa

Partially supported by Ronni Geraldo Gomes de Amorim

e-mail: ronniamorim@gmail.com

Faculdade do Gama

University of Brasília

Abstract

We will present a particular solution to the following D’Alambert non homogeneous partial differential equation

$$\frac{\partial^2 \psi(x, t)}{\partial x^2} - \frac{1}{v^2} \frac{\partial^2 \psi(x, t)}{\partial t^2} = \frac{K}{x^2 - (vt)^2}, \quad v \in \mathbb{R}^+$$

using a method based on complex analysis, adapted to another number set, the Perplex Numbers.

Positive ground states for a subcritical and critical coupled system involving Kirchhoff-Schrödinger equations

José Carlos de Albuquerque

Jointly work with Giovany M. Figueiredo (UnB) and João Marcos do Ó (UnB).

Research supported in part by INCTmat/MCT/Brazil

CNPq and CAPES/Brazil,

e-mail: joserre@gmail.com

Departament of Mathematics

University of Goiás

Goiânia, Brasil

Abstract

In this paper we prove the existence of positive ground state solution for a class of linearly coupled systems involving Kirchhoff-Schrödinger equations. We study the subcritical and critical case. Our approach is variational and based on minimization technique over the Nehari manifold. We also obtain a nonexistence result using a Pohozaev identity type.

References

- [1] C.O. ALVES, F.J.S.A. CORRÊA, AND T.F. MA, *Positive solutions for a quasilinear elliptic equation of Kirchhoff type*, Comput. Math. Appl. **49** (2005), 85-93.
- [2] C.O. ALVES, AND G.M. FIGUEIREDO, *Nonlinear perturbations of a periodic Kirchhoff equation in \mathbb{R}^N* , Nonlinear Anal. **75** (2012), 2750–2759.
- [3] J.M. DO Ó, AND J.C. DE ALBUQUERQUE, *Ground states for a linearly coupled system of Schrödinger equations on \mathbb{R}^N* , to appear in Asymptot. Anal.
- [4] H. SHI, AND H. CHEN, *Ground state solutions for asymptotically periodic coupled Kirchhoff-type systems with critical growth*, Math. Methods Appl. Sci. **39** (2016), 2193–2201.
- [5] X.H. TANG, AND S. CHEN, *Ground state solutions of Nehari-Pohozaev type for Kirchhoff-type problems with general potentials*, Calc. Var. Partial Differential Equations **56** (2017), 25pp.

Continuums of positive solutions for classes of non-autonomous and non-local problems with strong singular term

Carlos Alberto Santos
Partially supported by CAPES/Brazil
e-mail: csantos@unb.br
Departament of Mathematics
University of Brasília
Brasília, Brazil

Lais Santos
e-mail: matmslais@gmail.com
Department of Mathematics
University of Brasília
Brasília, Brazil

Pawan Kumar Mishra
Partially supported by CAPES/Brazil
e-mail: pawanmishra31284@gmail.com
Departament of Mathematics
University of Paraíba
João Pessoa, Brazil.

Abstract

In this work, we show existence of *continuums* of positive solutions for non-local quasilinear problems with strongly-singular reaction term on a bounded domain in \mathbb{R}^N , with $N \geq 2$. We approached non-autonomous and non-local equations by applying the Bifurcation Theory to the corresponding ϵ -perturbed problems and using a comparison principle for $W_{\text{loc}}^{1,p}(\Omega)$ -sub and supersolutions to obtain qualitative properties of the ϵ -*continuum* limit. Moreover, this technique empowers us to study a strongly-singular and non-homogeneous Kirchhoff problem to get the existence of a *continuum* of positive solutions.

References

- [1] Claudianor O. Alves and Dragoş-Pătru Covei. *Existence of solution for a class of nonlocal elliptic problem via sub-supersolution method*. Nonlinear Anal. Real World Appl., **23**:1–8, 2015.
- [2] David Arcoya, Tommaso Leonori and Ana Primo. *Existence of solutions for semilinear nonlocal elliptic problems via a Bolzano theorem*. Acta Appl. Math., **127**:87–104, 2013.
- [3] M. G. Crandall, P. H. Rabinowitz and L. Tartar. *On a Dirichlet problem with a singular non-linearity*. Comm. Partial Differential Equations, **2**(2):193–222, 1977.
- [4] Tarcyana S. Figueiredo-Sousa, Cristian Morales-Rodrigo and Antonio Suárez. *A non-local non-autonomous diffusion problem: linear and sublinear cases*. Z. Angew. Math. Phys., **68**(5):Art. 108, 20, 2017.
- [5] C. A. Santos and L. M. Santos. *How to break the uniqueness of $W_{\text{loc}}^{1,p}(\Omega)$ -solutions for very singular elliptic problems by non-local terms*. preprint available at <http://arxiv.org/abs/1805.10542>.

Singularity formation in toy models for incompressible flow

Leonardo Epiphany Galvão

Supported by Fapesp

e-mail: leonardo.epgalvao@gmail.com

Institute of Mathematics

Universidade Estadual de Campinas

Campinas, Brazil

Abstract

In this presentation, I will explore some toy models for the equations of incompressible fluid flow, in hopes of gaining insight on the possible devices for finite time singularity formation. This is achieved by examining geometric and/or analytic similarities between the models and the 3D Euler and Navier-Stokes equations. In doing so, I will survey through some of the classical techniques used in the theory of partial differential equations. More specifically, I deal with the well-understood CLM equation [2], alongside its proposed viscosity models [4, 3], in connection with the framework of 3D vortex stretching, and the equation discussed in [1], due to its similarities with the SQG equation, which itself can be viewed as a 2D model for the 3D Euler equations. These topics are part of my ongoing undergraduate research project funded by Fapesp, under the supervision of Prof. Anne Caroline Bronzi, in the process 2018/05899 – 5.

References

- [1] L. Silvestre, V. Vicol, *On a transport equation with nonlocal drift*, Trans. Amer. Math. Soc. 368, no. 9, 6159 – 6188, 2016.
- [2] P. Constantin, P. Lax, A. Majda, *A simple one-dimensional model for the three-dimensional vorticity equation*, Comm. Pure Appl. Math. 3, no. 6, 715 – 724, 1985.
- [3] T. Sakajo, *On global solutions for the Constantin-Lax-Majda equation with a generalized viscosity term*, Nonlinearity 16, no. 4, 1319 – 1328, 2003.
- [4] S. Schochet, *Explicit solutions of the viscous model vorticity equation*, Comm. Pure Appl. Math. 39, no. 4, 531 – 537, 1986.

A Brézis-Oswald problem to Φ -Laplacian operator with a gradient term

Marcos L. M. Carvalho

Jointly work with Goncalves, J. V., Silva, E. D.

Partially supported by CNPq, Santos, C. A.

Partially supported by PROCAD/Brazil

e-mail: marcos_leandro_carvalho@ufg.br

Instituto de Matemática e Estatística

Universidade Federal de Goiás

Goiânia - GO, Brazil

Abstract

It is established existence of minimal solution to the quasilinear elliptic problem

$$\begin{cases} -\Delta_{\Phi} u = \lambda f(x, u) + \mu g(x, u, \nabla u) & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \quad u = 0 & \text{on } \partial\Omega, \end{cases}$$

where f, g have a sublinear growth, $\lambda > 0$, and $\mu \geq 0$ are real parameters. Our results are an improvement of the classical Brézis-Oswald result to Orlicz-Sobolev setting by including singular nonlinearity as well as a gradient term.

References

- [1] Brézis, H. & Oswald, L., *Remarks on sublinear elliptic equations*, *Nonlinear Anal.* **10**, 55–64, 1986.
- [2] Carvalho, M. L. M., Goncalves, J. V., Silva, E. D., Santos, C. A., *A Type of Brézis-Oswald Problem to the Φ -Laplacian Operator with Very Singular Term*. *Milan J. Math.* **86**, 53–80, 2018.

Existence and uniqueness of solutions for RFDEs

Mateus Malta Fleury
Partially supported by CNPq
e-mail: mateus.fleury@hotmail.com
Departament of Mathematics
University of Brasilia
Brasilia, Brazil

Abstract

In this work, we define a retarded functional differential equation (RFDE, for short) given by:

$$\begin{cases} \dot{x}(t) = f(t, x_t) \\ x_\sigma = \phi. \end{cases}$$

with $\phi \in C([-r, 0], \mathbb{R}^n)$ and $x_t : [-r, 0] \rightarrow \mathbb{R}^n$ is such that $x_t(\theta) = x(t + \theta)$. We consider $x \in C([\sigma - r, \sigma + A], \mathbb{R}^n)$, $t \in [\sigma, \sigma + A]$, $\theta \in [-r, 0]$ and $f : \mathbb{R} \times C([-r, 0], \mathbb{R}^n) \rightarrow \mathbb{R}^n$. Also, we prove some results for this type of equation such as the existence and uniqueness of solutions.

References

- [1] J. K. Hale. *Introduction to functional differential equations*. Applied mathematical sciences, **447**, Springer, 1993.

Asymptotically Linear Indefinite Problems in \mathbb{R}^N via an Abstract Linking Theorem

Liliane A. Maia and Mayra S. C. Rodrigues

Partially supported by CNPQ

e-mails: lilimaia.unb@gmail.com ssc_mayra@hotmail.com

Department of Mathematics

University of Brasília

Brasília, Brazil.

Abstract

An abstract linking result for Cerami sequences is proved without assuming a compactness condition. It is applied directly in order to prove the existence of critical points for a class of indefinite problems in infinite dimensional Hilbert Spaces. The main applications are given to Hamiltonian systems and Schrödinger equations. Here spectral properties of the operators are exploited and hypotheses of monotonicity on the nonlinearities are discarded.

References

- [1] Benci, V. and Rabinowitz, P. H.: Critical Point Theorems for Indefinite Functionals. *Inventiones Math.* **52**, 241-273 (1979).

Existence of solutions in Besov-Morrey spaces for chemotaxis Navier-Stokes fluid with chemical attractant

Monisse Postigo Alves¹, Lucas C. F. Ferreira²

E-mail: alvesmonisse@gmail.com (M.P. Alves), lcff@ime.unicamp.br (L.C.F. Ferreira).

M.P. Alves was supported by CNPq, Brazil.

LCF Ferreira was supported by FAPESP and CNPq, Brazil.

Department of Mathematics

^{1,2} University of Campinas

Campinas, SP, Brazil.

Abstract

The goal of this project is to study the asymptotic behavior type to the Keller-Segel system coupled with Navier-Stokes fluid in \mathbb{R}^N ($N \geq 2$). We will present the result about existence of global mild solutions to these equations by taking initial data belonging to homogeneous Besov-Morrey spaces. The mild solutions are obtained by means of a fixed point argument in a time-dependent space X , where the space X is constructed so that its norm is invariant by the intrinsic scaling of the equations in question. These results are part of my PhD thesis.

References

- [1] H. Kozono, M. Miura, and Y. Sugiyama, Existence and uniqueness theorem on mild solutions to the Keller-Segel system coupled with the Navier-Stokes fluid, *Journal of Functional Analysis* 270(5) (2016), 1663-1683.
- [2] J. Zhao and J. Zhou, Temporal decay in negative Besov spaces for the 3D coupled chemotaxis-fluid equations, *Nonlinear Anal. Real World Appl.* 42 (2018), 160-179.

Global well-posedness of weak solutions for the Vlasov-Fokker-Planck system

Paulo N. Seminario Huertas

Partially supported by CAPES PROEX-8477445/D and by CNPq DS 141602/2018-0

e-mail: pseminar@icmc.usp.br

Institute of Mathematics and Computer Sciences

University of São Paulo

São Carlos-SP, Brazil

Abstract

This paper is devoted to show the existence of weak solutions of the kinetic Vlasov-Fokker-Planck system in bounded domains with the self-consistent force field bounded at any time. The existence and uniqueness of weak solutions is proved as in Carrillo [1]. This fact is analysed using a variational technique and the theory of elliptic-parabolic equations of second order.

References

- [1] J.A. Carrillo. *Global Weak Solutions for the Initial-Boundary-Value Problems to the Vlasov-Poisson-Fokker-Planck System*, Math. Meth. in the Appl. Sci. **21**, 907-938 (1998).
- [2] J.A. Carrillo and J. Soler. *On the initial value problem for the Vlasov-Poisson-Fokker-Planck system with initial data in L^p spaces*, Math. Meth. in the Appl. Sci., **18**, 825-839 (1995).
- [3] J. L. Lions. *Equations différentielles opérationnelles et problèmes aux limites*, Springer, Berlin, 1961.

On the Quantum Harmonic Oscillator Model applied to Spectroscopy

Pedro Gabriel Ferreira Colucci
Email: pedro.colucci@usp.br
Instituto de Física de São Carlos
Universidade de São Paulo, São Carlos, Brazil

Márcia Federson
Supported by FAPESP and CNPq
Email: federson@icmc.usp.br
Instituto de Ciências Matemáticas e de Computação
Universidade de São Paulo, São Carlos, Brazil

Abstract

One of the systems with great occurrence in nature is the Harmonic Oscillator. When it does not appear explicitly, one can use it to make an approximate analysis of how the system would behave at a certain limit. Here we analyze two aspects:

- how the molecules behave in infrared spectroscopy using the Quantum Oscillator, where the vibrational energies of the molecules can be estimated by solving the Schrödinger equation;
- how the electric field is used to generate dipoles, thus varying the absorption or the scattering of the infrared beam.

References

- [1] Mulati, A.C.N. *Avaliação Físico Química de complexos de incluso de insulina e curcumina em ciclodex-trinas: Estudo com as espectroscopias Raman, FTIR e Fotoacústica*. Tese de Doutorado, Universidade Estadual de Maringá, Maringá PR: 2015.

Existence of multiple positive solutions for Schrödinger equation with singular nonlinearities sign-changing

Carlos Alberto Santos

Partially supported by CAPES/Brazil

e-mail: csantos@unb.br

Departament of Mathematics

University of Brasília

Brasília, Brazil

Ricardo Lima Alves

Partially supported by CNPq/Brazil

e-mail: ricardoalveslima8@gmail.com

Departament of Mathematics

University of Brasília

Brasília, Brazil.

Kaye O. Silva

e-mail: kayeoliveira@hotmail.com

Department of Mathematics

University of Goiás

Goiânia, Brazil

Abstract

In this work, we study the Schrödinger equation

$$\begin{cases} -\Delta u + V(x)u = \lambda a(x)u^{-\gamma} + b(x)u^p & \text{in } \mathbb{R}^N \\ u > 0, & \text{in } \mathbb{R}^N, \end{cases}$$

where $\lambda > 0$ is a real parameter, $N \geq 3$, $0 < \gamma < 1 < p < 2^* - 1$, $a \in L^{\frac{2}{1+\gamma}}(\mathbb{R}^N)$, $b \in L^\infty(\mathbb{R}^N)$ and V is a potential satisfying $X \hookrightarrow L^q(\mathbb{R}^N)$, where $2 \leq q < 2^*$ and $X = \{u \in H^1(\mathbb{R}^N); \int V(x)u^2(x)dx < \infty\}$. The main novelties in our result are obtaining multiplicity of positive solutions beyond of the extremal that is classical to apply Nehari's methods before it and allowing the potential b changing its signal.

References

- [1] J. Graham-Eagle. *A variational approach to upper and lower solutions*. IMA J. Appl. Math., **44**: 181–184, 1990.
- [2] N. Hirano, C. Saccon, N. Shioji, *Existence of multiple positive solutions for singular elliptic problems with a concave and convex nonlinearities*. Adv. Differential Equations **9**: 197–220, 2004.
- [3] K. Silva, A. Macedo, *Local minimizers over the Nehari manifold for a class of concave-convex problems with sign changing nonlinearity*, J. Differential Equations, **265**(5), 1894–1921, 2018.

On Lane-Emden systems with singular nonlinearities

Rodrigo Clemente

Partially supported by CAPES/CNPq

e-mail: rodrigo.clemente@ufrpe.br

Department of Mathematics

Rural Federal University of Pernambuco

Recife, Brazil

João Marcos Bezerra do Ó

Partially supported by INCT-Mat, CAPES and CNPq

e-mail: jmbo@pq.cnpq.br

Department of Mathematics

Federal University of Paraíba

João Pessoa, Brazil

Abstract

We are interested in regularity properties of semi-stable solutions for a class of singular semi-linear elliptic problems with advection term defined on a smooth bounded domain of a complete Riemannian manifold with zero Dirichlet boundary condition. We prove uniform Lebesgue estimates and we determine the critical dimensions for these problems with nonlinearities of the type Gelfand, MEMS and power case. As an application, we show that extremal solutions are classical whenever the dimension of the manifold is below the critical dimension of the associated problem. Moreover, we analyze the branch of minimal solutions and we prove multiplicity results when the parameter is close to critical threshold and we obtain uniqueness on it. Furthermore, for the case of Riemannian models we study properties of radial symmetry and monotonicity for semi-stable solutions.

References

- [1] A. Ambrosetti, P. Rabinowitz: *Dual variational methods in critical point theory and applications*. J. Functional Analysis **14** (1973), 349–381.
- [2] P. Antonini, D. Mugnai, P. Pucci: *Quasilinear elliptic inequalities on complete riemannian manifold*. J. Math. Pures Appl. **87** (2007), 582–600.
- [3] E. Berchio, A. Ferrero, G. Grillo *Stability and qualitative properties of radial solutions of the Lane-Emden-Fowler equation on Riemannian models*. J. Math. Pure. Appl. **102**, 1–35, 2014.
- [4] H. Berestycki, A. Kiselev, A. Novikov, L. Ryzhik: *The explosion problem in a flow*. J. Anal. Math. **110** (2010), 31–65.
- [5] D. Castorina, M. Sanchón: *Regularity of stable solutions to semilinear elliptic equations on Riemannian models*. Adv. Nonlinear Anal. **4** (2015), 295–309.
- [6] X. Luo, D. Ye, F. Zhou: *Regularity of the extremal solution for some elliptic problems with singular nonlinearity and advection*. J. Differential Equations **251** (2011), 2082–2099.
- [7] M. do Carmo, C. Xia, *Complete manifolds with non-negative Ricci curvature and the Caffarelli-Kohn-Nirenberg inequalities*. Compos. Math. **140** (2004), 818–826.

A model for determining the concentration of urea in artificial kidney

Sanjeev Kumar
Department of Mathematics
School of Basic Sciences and Research Sharda University
Greater Noida , India

Alpna Mishra¹
e-mail: alpnamishra@gmail.com
Department of Mathematics
Dr. B.R. Ambedkar University
Agra, India

Abstract

In medicine, dialysis is primarily used to provide an artificial replacement for lost kidney function (renal replacement therapy) due to renal failure. Dialysis may be used for very sick patients, who have suddenly but temporarily lost their kidney function (acute renal failure) or for quite stable patients who have permanently lost their kidney function. We present a model, which consist of partial differential equation defining the process of diffusion in artificial kidney and finally get the solution of this equation in the form of concentration of urea in blood by using the finite difference approach. The graph drawn between concentration and radial distance shows the variation between these two quantities. Dialysis treatments replace some of these functions through diffusion (waste removal) and ultrafiltration (fluid removal).

Sistemas com Termo Cncavo-Convexo em Domnio No Limitado

Steffânio Moreno de Sousa
e-mail: steffaniom@gmail.com
Instituto de Matemática e Estatística
UFG, GO, Brasil

José Valdo Goncalves
e-mail: goncalves.jva@gmail.com
Instituto de Matemática e Estatística
UFG, GO, Brasil

Abstract

Este trabalho estabelece a existncia de soluo para Sistemas com Termo Cncavo-Convexo em Domnio No-Limitado para o operador p-Laplaciano. Estendendo, de certa forma, o problema estudado no artigo [3]. A grande dificuldade que surge ao considerar um sistema, a troca de informao para mostrar a existncia de uma super-soluo para o mesmo.

References

- [1] A. AMBROSETTI, H. BRÉZIS, G. CERAMI - Combined effects of concave and convex nonlinearities in some elliptic problems, *J. Func. Anal.* 122, **10**, (1994) 519-543.
- [2] CARRIAO, P. C. AND GONCALVES, J. V. AND MIYAGAKI, O. H. - *Existence and λ - behavior of positive solutions of the equation $-\Delta u + a(x)u = \lambda u^q + u^p$ in \mathbf{R}^N* , Comm. Appl. Nonlinear Anal., 1999.
- [3] CARRIÃO, P. C. AND GONCALVES, J. V. AND MIYAGAKI, O. H. - *Existence and nonexistence in a class of equations with supercritical growth*, Appl. Anal.,2000.

Fourth order elliptic problems with combined nonlinearities

Thiago Rodrigues Cavalcante
e-mail: thiago.cavalcante@uft.edu.br
Universidade Federal do Tocantins
Arraias-TO, Brazil

Abstract

It is establish existence and multiplicity of solutions for a class of superlinear elliptic involving a fourth order elliptic problem under Navier conditions on the boundary. Here we do not apply the well known Ambrosetti-Rabinowitz condition at infinity. Instead of we assume that the nonlinear term is a nonlinear function which is nonquadratic at infinity. The nonlinear term is a concave-superlinear function which is indefinite in sign. In order to apply variational methods we employ some fine arguments in order to recover the compactness

References

- [1] F.J.S.A. CORRÊA, J.V. GONÇALVES, A. RONCALLI, *On a Class of Fourth Order Nonlinear Elliptic Equations Under Navier Boundary Conditions*, Analysis and Applications, Anal. Appl. (Singap.) 8 (2010), 185-197.
- [2] J. V. A. GONÇALVES, *On bounded nonlinear perturbations of an elliptic equation at resonance* Nonlinear Anal., 5, (1981), 57-60.
- [3] J.V.A. GONÇALVES, E. D. SILVA, M. L. SILVA, *On positive solutions for a fourth order asymptotically linear elliptic equation under Navier boundary conditions*, 384, (2011), 387-399.
- [4] Y. YANG, J. H. ZHANG, *Existence of solutions for some fourth-order nonlinear elliptical equations*, J. Math. Anal. Appl. 351, (2009), 128-137.
- [5] Y. YE, C.L. TANG, *Existence and multiplicity of solutions for fourth-order elliptic equations in RN* , J. Math. Anal. Appl. 406 (2013), no. 1, 335-351.
- [6] DA SILVA, E. D. ,CAVALCANTE, T. R. *Multiplicity of solutions to fourth-order superlinear elliptic problems under Navier conditions*, Electronic Journal of Differential Equations. 167, (2017), 1-16.

Sponsors



FAP - DF
Fundação de Apoio à Pesquisa do Distrito Federal



UnB
Universidade de Brasília



FAPESP
Fundação de Amparo à Pesquisa do Estado de São Paulo



FINATEC
Fundação de Empreendimentos Científicos e Tecnológicos



Bento Viana

Index

- Adilson Presoto, 90
Agnieszka Malinowska, 38
Aldo Pereira, 31
Alex F. Pereira, 61
Alexandre Nolasco, 17
Alfredo Sotelo Pejerrey, 97
Anderson L.A. de Araujo, 91
André Caldas, 48
Angelina Carrijo, 98
Arlúcio Viana, 92
- Carlos Rocha, 18
Carlos Tomei, 39
Cláudia Santana, 100
Claudia Correa, 62
Claudiney Goulart, 101
- Damião J. Araújo, 75
Daniel Raom Santiago, 102
Daniela M. Vieira, 63
Diego Moreira, 76
- Eduardo Domingos, 77
Eduard Toon, 32
Elisa R. Santos, 64
Eudes Mendes Barboza, 80
Everaldo de Mello Bonotto, 49
- Felipe B. Federson, 103
Fernanda Andrade da Silva, 33
Fernando S. Carvalho, 104
- Gabriela Planas, 19
Gaetano Siciliano, 82
Gennaro Infante, 20
Geraldo Botelho, 66
Geraldo Silva, 34
Gustavo Araújo, 65
Gustavo Madeira, 81
- Hans-Otto Walther, 21
- Ian Rodrigues Massa, 105
- Jacson Simsen, 50
Jamil Abreu, 93
Jamilson R. Campos, 67
João Marcos do Ó, 40
Jorge Marques, 84
- José Carlos de Albuquerque, 106
José Galé, 22
José Francisco de Oliveira, 83
Juliana Berbert, 51
- Kaye Oliveira Silva, 85
Khazhak Navoyan, 68
- Lais Santos, 107
Leonardo Epiphany Galvão, 108
Lucas Seco, 52
Luis Henrique de Miranda, 94
- M. Lilian Lourenço, 69
Márcia Federson, 24
Mahamadi Warma, 23
Marccone C. Pereira, 54
Marcos L. M. Carvalho, 109
Maria Carolina Mesquita, 41
Marielle Silva, 42
Martin Bohner, 25
Mateus Malta Fleury, 110
Mauro Patrão, 53
Maxwell Lizete, 86
Mayra S. C. Rodrigues, 111
Minbo Yang, 87
Monisse Postigo Alves, 112
- Nilson C. Bernardes Jr., 70
- Olimpio H. Miyagaki, 26
- Pablo Amster, 43
Paulo N. Seminario Huertas, 113
Pawan Mishra, 88
Pedro Gabriel Ferreira Colucci, 114
Pedro J. Miana, 27
Phillipo Lappicy, 55
Pierluigi Benevieri, 44
Piotr Kalita, 56
- Ricardo Lima Alves, 115
Rodrigo A. Samprognna, 58
Rodrigo Clemente, 116
Rodrigo Nunes Monteiro, 95
- Sabrina Camargo, 59
Sabrina Streipert, 35
Sanjeev Kumar, 117

Santiago Muro, 71
Steffânio Moreno de Sousa, 118

Thiago Cavalcante, 119
Thiago R. Alves, 72
Tibor Krisztin, 28
Tom Cuchta, 36

Valeriano Antunes, 46
Victor Simões Barbosa, 73